

JAML, a novel protein with characteristics of a junctional adhesion molecule, is induced during differentiation of myeloid leukemia cells

Christel Moog-Lutz, Florence Cavé-Riant, Florence C. Guibal, Marie A. Breau, Yolande Di Gioia, Pierre Olivier Couraud, Yvon E. Cayre, Sandrine Bourdoulous, and Pierre G. Lutz

Retinoic acid induces clinical remission in acute promyelocytic leukemia (APL) by triggering differentiation of leukemia promyelocytes. Here, we have characterized a gene encoding a member of the immunoglobulin superfamily, among novel retinoic acid–induced genes identified in APL cells. This protein, which was named JAML (junctional adhesion molecule–like), contains 2 extracellular immunoglobulin-like domains, a transmembrane

segment, and a cytoplasmic tail. JAML mRNA is expressed in hematopoietic tissues and is prominently expressed in granulocytes. The fact that JAML protein is localized at the cell plasma membrane in the areas of cell-cell contacts, whereas it is not detected at free cell borders, suggests that JAML is engaged in homophilic interactions. Furthermore, a conserved dimerization motif among JAM members was shown to be important for

JAML localization at the cell membrane. Finally, exogenous expression of JAML in myeloid leukemia cells resulted in enhanced cell adhesion to endothelial cells. Altogether, our results point to JAML as a novel member of the JAM family expressed on leukocytes with a possible role in leukocyte transmigration. (Blood. 2003;102:3371-3378)

© 2003 by The American Society of Hematology

Introduction

Treatment of acute promyelocytic leukemia (APL) with retinoic acid (RA) is a seminal example of a therapeutic strategy aimed at targeting the activities of a specific oncoprotein with a cell differentiation agent.¹⁻³ The induction of remission is associated with the terminal differentiation of immature leukemic cells into neutrophils. In 95% of clinical APL cases, a specific translocation t(15;17)⁴ creates a promyelocytic leukemia–retinoic acid receptor alpha (PML-RAR α) fusion between the N-terminal region of the PML locus⁵⁻⁸ and the C-terminus of the RAR α .^{9,10} Expression of PML-RAR α in myeloid cell lines increases RA sensitivity¹¹ and RA-induced differentiation of APL cells is dependent on the presence of PML-RAR α . This suggests that differentiation of APL cells by RA is triggered through a PML-RAR α –dependent signaling pathway.

When hematopoietic cells leave the circulation, they tether to and roll on the endothelial cell surface, spread, and finally emigrate through the endothelial cells to reach the underlying tissues. Such mechanisms have been demonstrated for hematopoietic stem cell homing and engraftment¹² and for leukocytes emigration from the bloodstream in response to changes on the surface of blood vessels that signal injury or infection.^{13,14} Transient opening of interendothelial contacts allows leukocytes to emigrate through the endothelial cells.¹⁵ Although a key role has been demonstrated for platelet endothelial cell adhesion molecule (PECAM)¹⁶⁻¹⁸ and for junctional adhesion molecule 1 (JAM-1),¹⁹ the molecular mechanisms underlying the diapedesis of leukocytes through the endothelial monolayer are not as yet fully understood. PECAM and JAM share

structural and functional features. Both belong to the immunoglobulin superfamily (IgSF) and engage in homo- and heterotypic interactions.^{17,19} Two other members of the JAM family, JAM-2^{20,21} (also known as VE-JAM²²) and JAM-3,^{23,24} have been identified as components of interendothelial junctional complexes. On leukocytes, a JAM counter-receptor was postulated¹⁹ although it is still poorly characterized. JAM-3, which is up-regulated following T-cell activation, is such a JAM-2 counter-receptor.²³ More recently, JAM proteins were shown to interact with integrins. JAM-1 was shown to be a ligand of the β_2 integrin LFA-1 and to contribute to LFA-1–dependent transendothelial migration of T cells and neutrophils.²⁵ JAM-2 interacts with $\alpha_4 \beta_2$ integrins in T cells²⁶ and JAM-3 interacts with the β_2 integrin Mac-1 in platelets.²⁷

In order to identify RA-target genes, which are critical to the process of differentiation, we have used a differential screening strategy with the NB4 APL cell line, which harbors the t(15;17) translocation encoding PML-RAR α . These cells undergo granulocytic differentiation when treated with all-*trans* retinoic acid (ATRA).²⁸ Among 4 novel genes that we have cloned as induced by RA in NB4 cells, we have identified a junctional adhesion molecule–like gene here named JAML (junctional adhesion molecule–like). The predicted JAML protein contains a signal peptide, 2 IgSF domains, a transmembrane segment and an intracellular tail. JAML shares structural homologies with molecules involved in adhesion, such as JAM. Here we show that JAML is a novel JAM protein that is expressed in mature hematopoietic cells. Furthermore, JAML protein is localized at the cell plasma membrane in the

From Hôpital Robert Debré, Institut National de la Santé et de la Recherche Médicale (INSERM) Unité 417; and Institut Cochin, INSERM Unité 567, Centre National de la Recherche Scientifique (CNRS) UMR8104, Université Paris V, Paris, France.

Submitted November 14, 2002; accepted June 30, 2003. Prepublished online as *Blood* First Edition Paper, July 17, 2003; DOI 10.1182/blood-2002-11-3462.

Supported by INSERM, CNRS, and grants from the Association Pour la Recherche sur le Cancer, the Fondation de France, the Comité de Paris de la

Ligue Contre le Cancer, the Lady Tata Memorial Trust, and the Leukemia Research Foundation.

Reprints: Pierre G. Lutz, INSERM U417, Hôpital Robert Debré, 48 Boulevard Serurier, 75019 Paris, France; e-mail: lutz@rdebre.inserm.fr.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2003 by The American Society of Hematology

areas of cell-cell contacts whereas it was not detected at free cell borders, suggesting that JAML is engaged in homophilic interactions. Finally, JAML expression enhances myeloid leukemia cell adhesion to endothelial cells.

Materials and methods

Cell lines, human cells, culture conditions, induction, and measurements of differentiation

PLB-985,²⁹ NB4,²⁸ NB4.306,³⁰ and HL-60 cells³¹ were grown and used as described.^{32,33} HT-29 cells were grown on 9-cm Petri dishes in Dulbecco modified Eagle medium (DMEM) containing 4.5 g/L glucose and 5% fetal bovine serum (Invitrogen, Cergy Pontoise, France). COS-7 cells were grown in 10-cm Petri dishes in DMEM containing 1 g/L glucose, 5% fetal bovine serum, and 1X nonessential amino acids (Invitrogen). U937 and MDCK cells were grown in DMEM containing 10% fetal bovine serum. Human bone marrow endothelial cells (HBMECs³⁴; kindly provided by Dr B. Weksler, Weill Medical College of Cornell University, NY), were cultured in DMEM Glutamax supplemented with 10% heat inactivated fetal bovine serum, 7.5 µg/mL endothelial cell growth supplement (Sigma-Aldrich, St Louis, MO), 7 IU heparin (Sigma-Aldrich) and 10 mM HEPES. Cell viability was estimated using standard trypan blue dye exclusion assay. ATRA (Sigma-Aldrich) was dissolved in ethanol. Me₂SO (Sigma-Aldrich) was used at the final concentration of 1.25%. Phorbol 12-myristate 13-acetate (PMA; Sigma-Aldrich) in Me₂SO was used at a final concentration of 107 nM. The final concentration of Me₂SO was 0.002% and had no effect on differentiation. Differentiation was assessed by the percentage of cells with cell-associated nitro blue tetrazolium (NBT; Sigma-Aldrich), and cell morphology as examined under light microscopy on cytospin slides stained with May-Grünwald-Giemsa.

PML-RAR α -expressing primary APL cells obtained from patients diagnosed at Saint-Louis Hospital (Paris, France), which were purified and cultivated with 10⁻⁷ M ATRA as described,³⁵ were kindly provided by C. Chomienne. Mononuclear cells and granulocytes were isolated as described.^{32,33}

cDNA library construction and screening

RNAs from NB4 cells before and after 24-hour treatment with 9-*cis* RA, and HT-29 cells, were extracted using the guanidinium isothiocyanate lysis buffer and purified by centrifugation through cesium chloride cushion. Poly(A)⁺ RNA was purified using the Oligotex affinity resin (Qiagen, Hilden, Germany). An RA-treated NB4 cDNA library was constructed in the Uni-ZAP XR vector (Stratagene, La Jolla, CA) using the manufacturer's protocol. Using 5 µg poly(A)⁺ RNA, 3 million recombinant phages were obtained and 120 000 of them were directly plated and transferred onto nylon filter replicas (Hybond NX; Amersham Biosciences, Freiburg, Germany). Subtracted probes were prepared as previously described.³⁶ Briefly, the untreated NB4-specific probe (minus probe) was obtained from 2 µg NB4 poly(A)⁺ RNA that was reverse transcribed into 320 ng single-stranded cDNA and hybridized to 9.6 µg (30 \times excess) of HT-29 poly(A)⁺ RNA. The RA-treated NB4-specific probe (plus probe) was obtained from 3 µg RA-treated NB4 poly(A)⁺ RNA that was converted into 400 ng single-stranded cDNA and hybridized to 12 µg NB4 poly(A)⁺ RNA (30 \times excess). After hydroxyapatite chromatography, 10% and 8% of the cDNA remained single stranded for minus and plus probes, respectively. Both probes (6 ng) were labeled with α -[³²P] deoxycytidine triphosphate (dCTP) (111 TBq/mmol) using a random primer labeling kit (Rediprime; Amersham Biosciences) to obtain 4.8 \times 10⁷ and 7.6 \times 10⁷ cpm, respectively. Nylon replicas were first hybridized with the minus probe (0.8 \times 10⁶ cpm/mL) at 42°C in 50% formamide, 5 \times saline sodium citrate (SSC), 0.5% sodium dodecyl sulfate (SDS), 0.04% polyvinylpyrrolidone, 0.04% Ficoll, 20 mM sodium pyrophosphate pH 6.5, 10% dextran sulfate, and 100 µg/mL denatured salmon sperm DNA for 36 hours. Stringent washes were performed at 60°C in 0.2 \times SSC and 0.1% SDS. Filters were autoradiographed at -80°C for 12 hours and 144 hours

using BioMax MS films (Kodak, Rochester, NY). The minus probe was removed using alkali procedure, and filters were hybridized with the plus probe (1.3 \times 10⁶ cpm/mL) for 36 hours, stringently washed, and autoradiographed as described above. A secondary screening was performed using the same protocol on 885 plaques giving differential signals with the plus and minus probes.

Plasmid recovery and analysis

Pure plaques were directly recovered as bacterial colonies using the pBluescript/Lambda ZAPII in vivo excision system (Stratagene). Plasmid minipreparations were performed using the Biomek 2000 robot (Beckman Coulter, Brea, CA), digested with *EcoRI* and *XhoI*, loaded and electrophoresed into 2 parallel 1% agarose gels. Gels were then blotted onto Hybond-N nylon membranes (Amersham Biosciences), which were hybridized to the plus and minus probes. Inserts from differential clones were purified from agarose gels using spin-X columns (Corning, Corning, NY) and ³²P-labeled by random priming. These probes were hybridized to both replicas of the library and dot blots containing previously prepared plasmids in order to perform cross-hybridization analysis.

Sequencing and computer analysis

Plasmid DNA was purified through Nucleobond columns (Macherey-Nagel, Düren, Germany) and double-stranded DNA templates were sequenced using pBluescript-specific and internal primers. Sequence analysis and alignments were carried out with the GCG programs (Wisconsin Package, version 12.0-UNIX; Accelrys, Paris, France). Sequence homologies were identified using the FastA and BLAST programs by searching the GenBank (release 130.0) and EMBL (release 71.0) data banks and in the case of translated sequences by searching the SWISS-PROT (release 40.0), PIR/NBRF (release 73.0), and SPTREMBL (release 21.0) data banks. The *bos taurus* JAML protein was deduced by translation of accession number BG688339 in comparison to the human JAML sequence.

To determine a dendrogram with various IgSF family members containing 2 Ig domains, the alignment was performed using CLUSTALW.³⁷ The phylogenetic tree was constructed using the Neighbor-Joining/UPGMA method (version 3.6a3). The PROTDIST program (version 3.6) was used to compute distance matrix under the Jones, Taylor, and Thornton model of amino acids change. The phylogenetic tree was plotted using NJPlot.³⁸

JAML cDNA cloning

The JAML cDNA fragment isolated in the differential screening contained 1484 base pair (bp) corresponding to the 3' end of JAML cDNA. A remaining 5' cDNA sequence was amplified by rapid amplification of complementary DNA ends-polymerase chain reaction (RACE-PCR) from RA-treated NB4 RNA (with 5 \times 10⁻⁷ M for 48 hours) using the Marathon cDNA amplification kit (Clontech, Palo Alto, CA) with the following primer: 5'-TGGTGACAGAGTCCAGTCTATCTTG-3'. Using the same kit, a contiguous cDNA was generated by reverse transcriptase (RT)-PCR amplification of JAML from RA-treated NB4 RNA. To determine the chromosomal localization and the intron/exon structure of the JAML gene, its complete cDNA was used to retrieve genomic data from databases. Structures of the JAM-1, JAM-2, and JAM-3 have been previously described.²³

Eukaryotic expression vectors, in vivo expression, and protein extracts

The JAML coding sequence tagged with the hemagglutinin (HA) epitope at its C-terminus was subcloned into the p513 vector, a derivative of the pSG5 expression vector,³⁹ resulting in the p513-JAML-HA vector; and into the pMTCB6⁺-derived expression vector⁴⁰ under the control of the zinc-inducible sheep metallothioneine promoter resulting in the pMT-JAML-HA vector. Construction of the pMT-JAMLK54D-HA vector was achieved by using the QuickChange site-directed mutagenesis kit (Stratagene). The forward oligonucleotide sequence was used (mutated base pairs in bold):

5'-ACAGAAGACAAATGTATATTCGACATAGACTGGACTCTGTAC-3'. COS-7 cells were transfected using calcium phosphate coprecipitation⁴¹ of 0.5 μ g DNA vectors (adjusted to 14 μ g per 10-cm Petri dish with pBluescript carrier DNA). Medium was changed after 16 hours. Thirty-six hours after transfection, COS-7 cells were harvested, washed once in phosphate-buffered saline (PBS), and resuspended in Laemmli buffer. MDCK and U937 cells were washed twice in serum-free medium, once in Opti-MEM (Invitrogen), resuspended in the same medium at 15×10^6 cells in 0.5 mL, and electroporated (Gene Pulser; Biorad, Hercules, CA) at 300 V, 960 μ F with 20 μ g of the empty pMT, pMT-JAML-HA, and pMT-JAMLK54D-HA vectors. Cells were then cultured for 48 hours prior to selection with 0.4 μ g/mL G418 (Invitrogen).

Northern blot analysis

Total RNA extraction and hybridization were as described.³² Human RNA master blot and human immune system multiple tissue Northern blot II were obtained from Clontech. The JAML probe corresponded to the 3' end (1484 bp) of the JAML cDNA. Probes were labeled with α -[³²P] dCTP (111 TBq/mmol) using a random primer labeling kit (Amersham Bioscience). Radioactivity was detected and quantified using a Storm 860 Phosphor Imager (Amersham Bioscience).

Antibodies and Western blot analysis

A polyclonal rabbit antiserum was raised against the JAML coding sequence lacking N-terminal amino acids (1-19) fused with the glutathione S-transferase (GST) sequence generated by cloning the corresponding JAML sequence into a derivative of pGEX-3X (Amersham Biosciences). After 5 minutes boiling in Laemmli buffer, samples were resolved by SDS-polyacrylamide gel electrophoresis (PAGE), and transferred onto polyvinylidene difluoride (PVDF) membranes (Perkin-Elmer Life Sciences, Boston, MA). Background immunoreactivity was reduced by preincubating membranes with PBS containing 5% nonfat dry milk and 0.1% Tween 20 for 1 hour. Western analysis was carried out with anti-JAML diluted 1:2000 in PBS containing 0.1% Tween 20 or anti-HA monoclonal antibody (Cell Signaling Technology, Beverly, MA) diluted 1:10 000 in PBS containing 0.5% Tween 20. Detection of antibody binding was achieved with horseradish peroxidase-conjugated secondary antibodies (Jackson Laboratories, West Grove, PA). Enzymatic activity was detected using the chemiluminescence reagent plus kit (Perkin-Elmer Life Sciences) and autoradiography. Anti- β -actin antibody was purchased from Santa Cruz Biotechnology, Santa Cruz, CA.

In vitro transcription/translation

In vitro-translated proteins were synthesized with the p513-JAML-HA vector in the presence of [³⁵S]-methionine using the TNT Coupled Reticulocyte Lysate System (Promega, Madison, WI). Glycosylations were achieved using the canine microsomal membrane system as recommended by the manufacturer (Promega). Proteins were visualized by autoradiography after separation by SDS-PAGE.

Immunofluorescence

Transfected MDCK cells were grown on glass slides for 8 hours, and for zinc-induced expression of JAML, 125 μ M ZnSO₄ was added to the medium. Forty hours after zinc treatment, cells were washed and fixed for 5 minutes with 2% paraformaldehyde in PBS. Cells were permeabilized with PBS, 0.1% Triton X100 for 5 minutes. After a 30-minute treatment with blocking solution (PBS, 1% bovine serum albumin [BSA]), cells were stained for 1 hour with a 1:10 000 dilution of anti-HA monoclonal antibody. Goat anti-mouse-Cy3 (Jackson Laboratories) at a dilution of 1:400 was used as a secondary antibody. DNA was counterstained with Hoechst 33258 dye, and preparations were mounted in mowiol (Calbiochem, Novato, CA). Fluorescence was viewed using an Olympus microscope equipped with appropriate optics and filter module for Cy3 detection.

Adhesion assay

When confluent, HBMEC monolayers were either left untreated or pretreated with 100 U/mL recombinant tumor necrosis factor α (TNF α) for 24 hours in DMEM supplemented with 0.1% BSA. U937 cells were stably transfected with the pMT, pMT-JAML-HA, or pMT-JAMLK54D-HA expression vectors and either induced by addition of 100 μ M ZnSO₄ for 18 hours or left uninduced. U937 cell populations were fluorescently labeled by incubation for 1 hour in RPMI medium containing 10 μ L green CFMDA (Molecular Probes, Eugene, OR). After 3 washes in PBS, labeled U937 cells were resuspended in DMEM supplemented with 0.1% BSA and 10⁶ cells were added into each well of 24-well -plates containing confluent HBMEC monolayers. After 30 minutes of incubation at 37°C, nonadherent U937 cells were removed by 3 consecutive washes. Adherent U937 cells were then lysed with 1% Triton X100 and released fluorescence was quantified using a fluorometer (Fusion; Perkin-Elmer). The percentage of adherent U937 cells was calculated in each condition as the ratio between the mean fluorescence recovered in 4 replicate wells and the total amount of fluorescence corresponding to the labeled U937 cells incubated in each well.

Results

JAML mRNA is up-regulated in ATRA-treated APL cells

Using subtractive hybridization, we have identified 23 genes induced in RA-treated NB4 cells. Among 4 novel genes (288/JAML, 438, PRAM-1, and ASB-2; Figure 1), PRAM-1 encodes an adaptor protein functionally linked to an RA-dependent signaling pathway³³ and ASB-2, a protein involved in growth inhibition and commitment in myeloid leukemia cells.⁴² Nineteen other genes corresponded to known sequences in gene databases (Table 1). Most of the known sequences were characteristics of the myeloid lineage; 2 (*CAMPATH-1* and *GLUT5*) were not known to be induced by RA.

ATRA-induced granulocytic differentiation of NB4 cells was accompanied by progressive up-regulation of the JAML mRNA, which was detected 12 hours after treatment (Figure 2A). As expected, ATRA did not affect growth and differentiation (Figure 2Bi-ii, respectively) of NB4.306, an ATRA-resistant NB4 subclone, which has lost the capacity to encode an intact ATRA-binding PML-RAR α fusion protein.³⁰ In these cells, no induction of JAML mRNA was observed (Figure 2Biii). In contrast, expression of JAML mRNA was strongly up-regulated in NB4 cells induced to growth arrest and differentiation by ATRA (Figure 2B). JAML mRNA was also highly induced by ATRA in blasts freshly obtained from peripheral blood of newly diagnosed patients with APL (Figure 2C) and expressing the PML-RAR α gene. This




Figure 1. Identification of 4 novel genes induced in RA-treated NB4 cells. (A) Autoradiogram of 288/JAML, 438, 809, and 813 mRNA expression in NB4 cells either untreated or treated for 24 hours with 5×10^{-7} M 9-cis RA, and in HT-29 cells as a nonhematopoietic negative control. Positions of the 28S and 18S rRNAs are indicated on the left side. The lower part is methylene blue-stained 28S and 18S rRNAs on membranes after transfer as assessment of RNA quantities in each lane (5 μ g total RNA).

Table 1. Characteristics of the differential cDNAs identified

Clone	Description	GenBank accession no.
3	5-lipoxygenase activating protein (FLAP)	X52195
13	Monocyte chemoattractant protein 1 (MCP-1)	X14768
16	Neutrophil cytosol factor 1 (p47phox)	M25665
20	GOS2	M69199
26	Cellular transglutaminase homologue (TGase H)	M98479
301	CD11b	J03925
617	Plasminogen activator inhibitor-2 (PAI-2)	J03603
685	Lymph node homing receptor (L-selectin)	M25280, X16070
685	Leu-8	X17519
685	LAM-1	X16150
732	Interleukin-8 (IL-8)	M17017
748	Bfl-1/A1	U27467, U29680
752	Defensin 1	M26602, M21130
756	Corticostatin HP-4 (defensin 4)	X65977
781	Delta-aminolevulinic synthase (ALAS1)	X56351
796	CAMPATH-1 (CD52)	X62466
806	Coactosin-like protein (CLP)	L54057
808	ICAM-1	J03132
820	c-fgr	M19722
832	GLUT5 glucose transporter	M55531
850	Src-like adapter protein (SLAP)	D89077

Leu-8 indicates leukocyte surface antigen Leu-8; LAM-1, leukocyte adhesion molecule-1; Bfl-1/A1, Bcl-2-related gene expression in fetal liver-1/Bcl-2-related protein A1; ICAM-1, intercellular adhesion molecule-1; and c-fgr, Gardner-Rasheed feline sarcoma viral proto-oncogene.

induction correlated with ATRA-induced differentiation of these cells as assessed morphologically and by NBT reduction assay (data not shown). Altogether, our data strongly suggested that increased JAML mRNA expression was associated with the capacity of APL cells to undergo granulocytic differentiation when treated with ATRA.

The JAML protein is a novel member of the IgSF localized at areas of cell-cell contacts

A 1638-bp JAML cDNA (GenBank Accession no. AJ515553) obtained from ATRA-treated NB4 cells (Figure 3A) contains a complete open reading frame predicted to encode a protein of 394 amino acids followed by a 3'-untranslated region and a poly(A) tail. A 19 amino acids signal peptide was predicted using SignalP.⁴³ Overall, the cDNA is predicted to encode a type I integral transmembrane protein with an extracellular portion of 275 amino acids, a transmembrane segment of 21 amino acids, and a cytoplasmic tail of 98 amino acids. The extracellular region of JAML comprises 2 Ig domains resembling surface antigens such as epithelial v-like antigen (EVA),⁴⁴ human Coxsackie and adenovirus receptor (HCAR),^{45,46} and JAM proteins.²³ The ectodomain pos-

sesses 2 Ig-like folds, the N-terminal domain is a C2-type whereas the membrane proximal domain is a V-type. When compared with other IgSF members, JAML appears to be related to the JAM family (Figure 3B). A motif (KID58 flanked by Y72 in human, KVD58 flanked by F72 in bos taurus) similar to the dimerization motif R(V, I, L)E flanked by an aromatic amino acid Tyr/Phe which was shown to be essential for JAM dimers formation in solution⁴⁷ was found in JAML proteins (Figure 3C). By using information derived from the chromosome 11 sequence, we mapped the human JAML gene to the 11q23 region. A remarkable conservation of gene structure and phase of introns within the members of the JAM family points to their evolutionary relationship (Table 2). Divergence between JAML and JAM-1, JAM-2, and JAM-3 occurs in the 5' noncoding region of JAML, which is encoded by a separate exon, and in the first IgSF domain, which is encoded by 2 exons. Altogether, this indicates that we have identified a novel protein exhibiting homologies with members of the JAM family.

In COS cells transfected with JAML in frame with a carboxy-terminal HA epitope tag, both anti-JAML polyclonal serum and anti-HA monoclonal antibodies detected 3 bands with molecular weights of 60 kDa, 58 kDa, and 50 kDa. We interpreted these bands as corresponding to different glycoforms of the protein (Figure 4Ai-ii). Indeed, when microsomal membranes were added to an in vitro-coupled transcription/translation reaction, a higher-molecular-weight species was detected corresponding to a glycosylated JAML protein (Figure 4Aiii). In ATRA-treated NB4 cells, anti-JAML polyclonal antibodies detected a specific 60 kDa protein 48 hours after treatment (Figure 4B). To investigate the cellular distribution of the JAML protein, a JAML expression vector (pMT-JAML-HA) under the control of the zinc-inducible metallothionein promoter was stably transfected into MDCK cells. As shown in Figure 4Ciii, the JAML protein was present at the cell plasma membrane only in areas of cell-cell contacts and remained diffuse or absent at free cell borders. Mutation of the dimerization motif by substitution of lysine 54 by aspartic acid abolished membrane localization of the JAMLK54D mutated protein (Figure 4Cvii). Altogether, our results indicated that the JAML protein was localized at the cell plasma membrane and suggested that it may engage in homophilic interactions via its dimerization interface.

JAML mRNA is up-regulated during induced differentiation of myeloid leukemia cells and is expressed in normal hematopoietic tissues

In HL-60 and PLB-985 cells, JAML mRNA was progressively expressed as cells were induced to differentiation toward the granulocytic pathway with RA or Me₂SO, and toward the monocytic pathway with PMA (Figure 5A). JAML mRNA was mainly





Figure 2. Induction of JAML mRNA in RA-treated cells. (A) Time-response to ATRA in NB4 cells. Cells were treated for different times using 5×10^{-7} M ATRA. (B) JAML expression correlates with the capacity of NB4 cells to differentiate. NB4 and NB4.306 cells were cultured with 10^{-6} M ATRA and harvested after 0, 24, and 48 hours. Viable cells (i) and percent of NBT-positive cells (ii) were counted each day and JAML mRNA expression was analyzed (iii). (C) JAML mRNA is induced in ATRA-treated primary APL cells. Cells were purified from 3 untreated patients cultivated in the absence (-) or presence (+) of 10^{-7} M ATRA for 5 (APL no. 2) or 6 (APL no. 1 and APL no. 3) days. Northern blots were performed using 8 μ g (A,Biii) or 2 μ g (C) total RNA. Hybridization with a glyceraldehyde phosphate dehydrogenase (GAPDH) probe controlled for RNA quantities in each lane.

Figure 3. JAML is related to JAM proteins. (A) cDNA and deduced amino acid sequences of the human JAML. Nucleotide residues are numbered in the 5' to 3' orientation, and amino acids in the reading frame are designated by the one-letter code. Asterisk indicates the stop codon. The putative hydrophobic signal peptide (underlined) and transmembrane sequences (dotted underline) are marked. Potential N-linked glycosylation sites are shaded. Cysteines likely to form disulfide bonds in the 2 Ig domains are circled. (B) Dendrogram of various IgSF family members containing 2 Ig domains (JAM-1, JAM-2, JAM-3, JAML, Coxsackie and adenovirus receptor (CAR), and endothelial cell-selective adhesion molecule (ESAM)). (C) Alignment of the amino-terminal region of the membrane distal Ig domains of these proteins and their mouse (JAM-1, JAM-2, JAM-3, CAR, ESAM) or bos taurus (JAML) homologs. Shaded regions represent residues identical between at least 4 proteins. The dimerization motif is boxed.



expressed in peripheral blood leukocytes and to a lower extent in bone marrow, spleen, and lymph nodes (Figure 5Bi). In freshly purified mononuclear cells, JAML mRNA was expressed in human granulocytes as well as, albeit to a much lesser extent, in monocytes and lymphocytes (Figure 5Bii). Using the Human RNA Master Blot, JAML mRNA was also expressed in fetal liver, fetal

spleen, and fetal thymus but was not detected in nonhematopoietic tissues, except lung which is known to contain a number of myelomonocytic cells (data not shown). Altogether, our results indicated that JAML mRNA was expressed during normal myelopoiesis as well as in leukemia cells induced to myeloid differentiation.

Table 2. Comparison of gene structure between JAM family members

HGNC exon	No.	JAM-1 1q21.2-21.3	<i>i</i>	JAM-2 21q21.2	<i>i</i>	JAM-3 11q25	<i>i</i>	JAML 11q23.3	<i>i</i>
5'	0	—	—	—	—	—	—	>73	—
Intron	0-1	—	—	—	—	—	—	10 054	—
ATG/signal sequence	1	103	—	>301	—	>76	—	64	—
Intron	1-2	44 222	1	43 994	1	79 105	1	2262	1
EC, Ig-fold 1	2	59	—	66	—	66	—	—	—
Intron	2-3	167	1	5916	1	740	1	—	—
EC, Ig-fold 1	3	98	—	108	—	114	—	153	—
Intron	3-4	252	1	3782	1	3470	1	1694	1
EC, Ig-fold 1	4	137	—	153	—	153	—	228	—
Intron	4-5	282	1	4768	1	398	1	4493	1
EC, Ig-fold 2	5	203	—	203	—	203	—	110	—
Intron	5-6	167	0	3290	0	951	0	2217	0
EC, Ig-fold 2	6	103	—	100	—	100	—	238	—
Intron	6-7	128	1	3709	1	2511	1	2815	1
Transmembrane	7	108	—	108	—	130	—	137	—
Intron	7-8	231	1	3347	1	—	—	2384	1
IC	8	13	—	16	—	—	—	95	—
Intron	8-9	304	2	2890	2	87	2	1176	2
IC	9	49	—	43	—	55	—	88	—
Intron	9-10	136	0	2257	0	321	0	2318	0
IC/STOP	10	>937	—	>283	—	>36	—	>452	—

HGNC indicates HUGO Nomenclature Committee; *i*, phase of introns; EC, extracellular domain; IC, intracellular domain; and Ig-fold, immunoglobulin-like fold.




Figure 4. JAML is a transmembrane protein induced upon ATRA-treatment in NB4 cells. (A) COS-7 cells were mock-transfected (-) or transfected with the p513-JAML-HA vector (JAML). Cell lysates were separated by SDS-PAGE and subjected to immunoblotting with anti-JAML (i) or anti-HA (ii) antibodies. The arrowhead indicates the major specific band. Brackets indicate minor specific bands likely corresponding to glycosylated JAML protein. (iii) JAML protein was in vitro-translated in the absence (-) or presence (+) of canine microsomal membranes. Arrow indicates a glycosylated JAML protein. (B) JAML protein expression during ATRA-induced differentiation of NB4 cells. NB4 cells were treated with 5×10^{-7} M ATRA for different times. Protein lysates corresponding to 10^5 cells were analyzed by Western blot, using anti-JAML or anti- β -actin polyclonal antibodies. In panels A and B, asterisks indicate nonspecific bands. (C) Cellular localization of JAML. MDCK cells stably transfected with the pMT-JAML-HA or pMT-JAMLK54D-HA were either untreated (-Zn) or treated (+Zn) with $125 \mu\text{M}$ ZnSO_4 for 40 hours. Cells were treated with anti-HA monoclonal antibody and stained with Cy3-conjugated goat anti-mouse antibody (ii,iii,vi, vii). In panels i, iv, v, and viii, the same cells as in panels ii, iii, vi, and vii, respectively, were counterstained with Hoechst 33258 reagent.

JAML facilitates leukocyte adhesion to endothelial cells

To establish whether JAML can facilitate the interaction between leukocytes and endothelial cells, U937 cells were stably transfected with the pMT-JAML-HA or pMT-JAMLK54D-HA constructs, or with the empty vector as control. These nonclonal cell lines were fluorescently labeled and subjected to a cell adhesion assay using HBMECs as a target layer, either treated by $\text{TNF}\alpha$ or not treated. As expected, only a low level of control U937 cell adhesion to endothelial cells was observed and this adhesion was largely

enhanced by treatment of the endothelial cell monolayer with $\text{TNF}\alpha$ (Table 3). We observed that JAML expression induced significantly both basal and cytokine-induced U937 cell adhesion to endothelial cell monolayers, whereas the expression of JAMLK54D mutated protein had no effect (Table 3). These results strongly suggest that JAML expression enhances myeloid leukemia cell adhesion to endothelial cells.

Discussion

We have identified a novel protein, JAML, exhibiting homologies with members of the JAM family. JAML was induced by ATRA in human myeloid leukemia promyelocytes. JAML mRNA was up-regulated in myeloid leukemia cells induced to differentiate




Figure 5. JAML mRNA is up-regulated during induced differentiation of myeloid leukemia cells and is expressed in normal hematopoietic tissues. (A) Up-regulation of JAML mRNA during induced differentiation of myeloid leukemia cells. Northern blots were performed using $5 \mu\text{g}$ total RNA. Hybridization with a GAPDH probe controlled for RNA quantities in each lane. (B) Autoradiograms of JAML mRNA expression in immune tissues (i) and peripheral blood cells (ii). Northern blots were either obtained from Clontech (i) or were performed using $5 \mu\text{g}$ total RNA from granulocytes, monocytes, and lymphocytes (ii). In panels A and B, GAPDH was used as a probe for assessment of RNA quantities in each lane.

Table 3. JAML expression enhances U937 cell adhesion to HBMECs

	HBMECs	HBMECs + $\text{TNF}\alpha$
U937/MT		
- Zn	1.00 \pm 0.36	5.29 \pm 0.17
+ Zn	1.07 \pm 0.33	4.79 \pm 0.29
U937/MT-JAML		
- Zn	1.00 \pm 0.30	6.74 \pm 0.52
+ Zn	2.65 \pm 0.17	11.04 \pm 1.39
U937/MT-JAMLK54D		
- Zn	1.00 \pm 0.13	4.83 \pm 0.30
+ Zn	1.38 \pm 0.47	4.23 \pm 0.47

U937/MT, U937/MT-JAML-HA, and U937/MT-JAMLK54D-HA cells stably transfected with the pMT, the pMT-JAML-HA, or pMT-JAMLK54D-HA respectively were untreated (-Zn) or treated (+Zn) with ZnSO_4 for 18 hours prior to fluorescent labeling and subjected to a cell adhesion assay using human bone marrow endothelial cells either untreated (HBMECs) or treated (HBMECs + $\text{TNF}\alpha$) as a target layer. Results are expressed in each condition as the amount of released fluorescence from adherent cells, normalized to unity for the corresponding -Zn/- $\text{TNF}\alpha$ condition. The adhesion level of U937/MT-JAML-HA cells onto untreated or $\text{TNF}\alpha$ -pretreated HBMECs, in the absence of Zn treatment (-Zn), was measured as 2.3% and 15.5% respectively, of the total number of incubated cells. Values shown are \pm SD.

through the granulocytic and monocytic pathways and was expressed in normal hematopoietic cells. JAML is structurally similar to transmembrane proteins involved in cell-cell contacts and was localized to cell-cell borders. Induced expression of an exogenous JAML in myeloid leukemia cells resulted in enhanced cell adhesion to endothelial cells.

JAML is predicted to encode a type I integral transmembrane protein with 2 extracellular IgSF domains characteristic of proteins implicated as mediators of cell-cell interactions, generally through heterophilic interactions with other IgSF members. Searches of DNA and protein databases revealed that the JAML sequence has homologies with Ig domains of HCAR,^{45,46} EVA,⁴⁴ and myelin P0.⁴⁸ Furthermore, JAML shares structural features with members of the JAM family.¹⁹⁻²⁴ We mapped the human JAML gene to the 11q23 region; whereas JAM-1 has been found at 1q21 and JAM-2 at 21q21. Although, the JAM genes are not clustered, the JAML gene lies upstream of the ESAM (endothelium cell-selective adhesion molecule, 11q24), CTH genes (11q24) and JAM-3 (11q25).²³ Interestingly, 2 salt bridges formed in a complementary manner by a novel dimerization motif were shown to be essential for JAM dimers formation in solution.⁴⁷ A similar motif in the JAML protein was shown to be required for JAML localization at the plasma membrane, suggesting that this protein may engage in homophilic interactions via this dimerization interface. This is further emphasized by the fact that JAML protein is localized at the cell plasma membrane in the areas of cell-cell contacts, whereas it was not detected at free cell borders.

We have found that the drastic increase in JAML mRNA expression induced by RA in NB4 cells correlated with growth

arrest and differentiation of these cells. Cycloheximide treatment did block RA induction of JAML mRNA (data not shown), suggesting that its expression was not regulated at the transcriptional level. JAML mRNA increased during RA- or Me₂SO-induced granulocytic differentiation and by PMA, which induced monocytic differentiation of both HL-60 and PLB-985 cells. Consistent with JAML expression in myeloid leukemia cells induced to differentiate, JAML mRNA was expressed in normal granulocytes and monocytes. Altogether, our results point to JAML as a novel member of the JAM family expressed on leukocytes. JAM family members are modulators of leukocyte adhesion to and transmigration through endothelial monolayers. Indeed, we have demonstrated here that JAML can facilitate the adhesion of U937 cells to endothelial cells. Furthermore, the observation that a JAML dimerization mutant protein has no effect strongly suggests that JAML may support adhesion of leukocytes to endothelial cells by engaging either in homotypic or heterotypic interactions with endothelial cell adhesion partners. Whether JAML supports leukocyte transmigration and possibly interacts with viral protein to mediate viral internalization as described for JAM-1⁴⁹ and HCAR^{45,46} will be further investigated.

Acknowledgments

We thank Drs A. Benecke and R. M. Mège for helpful comments and suggestions and Dr B. Weksler for kindly providing us with the HBMECs. Sequencing was conducted by C. Cruaud at the Centre National de Séquençage (Evry, France).

References

- Huang ME, Ye YC, Chen SR, et al. Use of all-trans retinoic acid in the treatment of acute promyelocytic leukemia. *Blood*. 1988;72:567-572.
- Castaigne S, Chomienne C, Daniel MT, et al. All-trans retinoic acid as a differentiation therapy for acute promyelocytic leukemia, I: clinical results. *Blood*. 1990;76:1704-1709.
- Warrell RP Jr, Frankel SR, Miller WH, et al. Differentiation therapy of acute promyelocytic leukemia with tretinoin (all-trans-retinoic acid). *N Engl J Med*. 1991;324:1385-1393.
- Rowley JD, Golomb HM, Dougherty C. 15/17 translocation, a consistent chromosomal change in acute promyelocytic leukaemia. *Lancet*. 1977;1:549-550.
- de The H, Lavau C, Marchio A, Chomienne C, Degos L, Dejean A. The PML-RAR alpha fusion mRNA generated by the t(15;17) translocation in acute promyelocytic leukemia encodes a functionally altered RAR. *Cell*. 1991;66:675-684.
- Goddard AD, Borrow J, Freemont PS, Solomon E. Characterization of a zinc finger gene disrupted by the t(15;17) in acute promyelocytic leukemia. *Science*. 1991;254:1371-1374.
- Kakizuka A, Miller WH Jr, Umehara K, et al. Chromosomal translocation t(15;17) in human acute promyelocytic leukemia fuses RAR alpha with a novel putative transcription factor, PML. *Cell*. 1991;66:663-674.
- Pandolfi PP, Grignani F, Alcalay M, et al. Structure and origin of the acute promyelocytic leukemia myl/RAR alpha cDNA and characterization of its retinoid-binding and transactivation properties. *Oncogene*. 1991;6:1285-1292.
- Borrow J, Goddard AD, Sheer D, Solomon E. Molecular analysis of acute promyelocytic leukemia breakpoint cluster region on chromosome 17. *Science*. 1990;249:1577-1580.
- de The H, Vivanco-Ruiz MM, Tiollais P, Stunnenberg H, Dejean A. Identification of a retinoic acid responsive element in the retinoic acid receptor beta gene. *Nature*. 1990;343:177-180.
- Grignani F, Ferrucci PF, Testa U, et al. The acute promyelocytic leukemia-specific PML-RAR alpha fusion protein inhibits differentiation and promotes survival of myeloid precursor cells. *Cell*. 1993;74:423-431.
- Peled A, Kollet O, Ponomarev T, et al. The chemokine SDF-1 activates the integrins LFA-1, VLA-4, and VLA-5 on immature human CD34(+) cells: role in transendothelial/stromal migration and engraftment of NOD/SCID mice. *Blood*. 2000;95:3289-3296.
- Springer TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell*. 1994;76:301-314.
- Cines DB, Pollak ES, Buck CA, et al. Endothelial cells in physiology and in the pathophysiology of vascular disorders. *Blood*. 1998;91:3527-3561.
- Dejana E, Corada M, Lampugnani MG. Endothelial cell-to-cell junctions. *Faseb J*. 1995;9:910-918.
- Vaporciyan AA, DeLisser HM, Yan HC, et al. Involvement of platelet-endothelial cell adhesion molecule-1 in neutrophil recruitment in vivo. *Science*. 1993;262:1580-1582.
- Muller WA, Weigl SA, Deng X, Phillips DM. PECAM-1 is required for transendothelial migration of leukocytes. *J Exp Med*. 1993;178:449-460.
- Bogen S, Pak J, Garifallou M, Deng X, Muller WA. Monoclonal antibody to murine PECAM-1 (CD31) blocks acute inflammation in vivo. *J Exp Med*. 1994;179:1059-1064.
- Martin-Padura I, Lostaglio S, Schneemann M, et al. Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J Cell Biol*. 1998;142:117-127.
- Cunningham SA, Arrate MP, Rodriguez JM, et al. A novel protein with homology to the junctional adhesion molecule: characterization of leukocyte interactions. *J Biol Chem*. 2000;275:34750-34756.
- Aurrand-Lions MA, Duncan L, Du Pasquier L, Imhof BA. Cloning of JAM-2 and JAM-3: an emerging junctional adhesion molecular family? *Curr Top Microbiol Immunol*. 2000;251:91-98.
- Palmeri D, van Zante A, Huang CC, Hemmerich S, Rosen SD. Vascular endothelial junction-associated molecule, a novel member of the immunoglobulin superfamily, is localized to intercellular boundaries of endothelial cells. *J Biol Chem*. 2000;275:19139-19145.
- Arrate MP, Rodriguez JM, Tran TM, Brock TA, Cunningham SA. Cloning of human junctional adhesion molecule 3 (JAM3) and its identification as the JAM2 counter-receptor. *J Biol Chem*. 2001;276:45826-45832.
- Aurrand-Lions M, Duncan L, Ballestrem C, Imhof BA. JAM-2, a novel immunoglobulin superfamily molecule, expressed by endothelial and lymphatic cells. *J Biol Chem*. 2001;276:2733-2741.
- Ostermann G, Weber KS, Zernecke A, Schroder A, Weber C. JAM-1 is a ligand of the beta(2) integrin LFA-1 involved in transendothelial migration of leukocytes. *Nat Immunol*. 2002;3:151-158.
- Cunningham SA, Rodriguez JM, Arrate MP, Tran TM, Brock TA. JAM2 interacts with alpha4beta1. Facilitation by JAM3. *J Biol Chem*. 2002;277:27589-27592.
- Santoso S, Sachs UJ, Kroll H, et al. The junctional adhesion molecule 3 (JAM-3) on human platelets is a counterreceptor for the leukocyte integrin Mac-1. *J Exp Med*. 2002;196:679-691.
- Lanotte M, Martin-Thouvenin V, Najman S, Balerini P, Valensi F, Berger R. NB4, a maturation inducible cell line with t(15;17) marker isolated

- from a human acute promyelocytic leukemia (M3). *Blood*. 1991;77:1080-1086.
29. Tucker KA, Lilly MB, Heck L Jr, Rado TA. Characterization of a new human diploid myeloid leukemia cell line (PLB-985) with granulocytic and monocytic differentiating capacity. *Blood*. 1987;70:372-378.
 30. Dermime S, Grignani F, Clerici M, et al. Occurrence of resistance to retinoic acid in the acute promyelocytic leukemia cell line NB4 is associated with altered expression of the pml/RAR alpha protein. *Blood*. 1993;82:1573-1577.
 31. Gallagher R, Collins S, Trujillo J, et al. Characterization of the continuous, differentiating myeloid cell line (HL-60) from a patient with acute promyelocytic leukemia. *Blood*. 1979;54:713-733.
 32. Lutz PG, Houzel-Charavel A, Moog-Lutz C, Cayre YE. Myeloblastin is an Myb target gene: mechanisms of regulation in myeloid leukemia cells growth-arrested by retinoic acid. *Blood*. 2001;97:2449-2456.
 33. Moog-Lutz C, Peterson EJ, Lutz PG, et al. PRAM-1 is a novel adaptor protein regulated by retinoic acid (RA) and promyelocytic leukemia (PML)-RA receptor alpha in acute promyelocytic leukemia cells. *J Biol Chem*. 2001;276:22375-22381.
 34. Schweitzer KM, Vicart P, Delouis C, et al. Characterization of a newly established human bone marrow endothelial cell line: distinct adhesive properties for hematopoietic progenitors compared with human umbilical vein endothelial cells. *Lab Invest*. 1997;76:25-36.
 35. Chomienne C, Ballerini P, Balitrand N, et al. All-trans retinoic acid in acute promyelocytic leukemias, II: in vitro studies: structure-function relationship. *Blood*. 1990;76:1710-1717.
 36. Tomasetto C, Regnier C, Moog-Lutz C, et al. Identification of four novel human genes amplified and overexpressed in breast carcinoma and localized to the q11-q21.3 region of chromosome 17. *Genomics*. 1995;28:367-376.
 37. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res*. 1994;22:4673-4680.
 38. Perriere G, Gouy M. WWW-query: an on-line retrieval system for biological sequence banks. *Biochimie*. 1996;78:364-369.
 39. Green S, Issemann I, Sheer E. A versatile in vivo and in vitro eukaryotic expression vector for protein engineering. *Nucleic Acids Res*. 1988;16:369.
 40. Cook DM, Hinkes MT, Bernfield M, Rauscher FJ III. Transcriptional activation of the syndecan-1 promoter by the Wilms' tumor protein WT1. *Oncogene*. 1996;13:1789-1799.
 41. Chen C, Okayama H. High-efficiency transformation of mammalian cells by plasmid DNA. *Mol Cell Biol*. 1987;7:2745-2752.
 42. Guibal FC, Moog-Lutz C, Smolewski P, et al. ASB-2 inhibits growth and promotes commitment in myeloid leukemia cells. *J Biol Chem*. 2002;277:218-224.
 43. Nielsen H, Engelbrecht J, Brunak S, von Heijne G. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng*. 1997;10:1-6.
 44. Guttinger M, Sutti F, Panigada M, et al. Epithelial V-like antigen (EVA), a novel member of the immunoglobulin superfamily, expressed in embryonic epithelia with a potential role as homotypic adhesion molecule in thymus histogenesis. *J Cell Biol*. 1998;141:1061-1071.
 45. Bergelson JM, Cunningham JA, Droguett G, et al. Isolation of a common receptor for Coxsackie B viruses and adenoviruses 2 and 5. *Science*. 1997;275:1320-1323.
 46. Tomko RP, Xu R, Philipson L. HCAR and MCAR: the human and mouse cellular receptors for subgroup C adenoviruses and group B coxsackieviruses. *Proc Natl Acad Sci U S A*. 1997;94:3352-3356.
 47. Kostrewa D, Brockhaus M, D'Arcy A, et al. X-ray structure of junctional adhesion molecule: structural basis for homophilic adhesion via a novel dimerization motif. *Embo J*. 2001;20:4391-4398.
 48. Lemke G, Axel R. Isolation and sequence of a cDNA encoding the major structural protein of peripheral myelin. *Cell*. 1985;40:501-508.
 49. Barton ES, Forrest JC, Connolly JL, et al. Junction adhesion molecule is a receptor for reovirus. *Cell*. 2001;104:441-451.



blood[®]

2003 102: 3371-3378

doi:10.1182/blood-2002-11-3462 originally published online
July 17, 2003

JAML, a novel protein with characteristics of a junctional adhesion molecule, is induced during differentiation of myeloid leukemia cells

Christel Moog-Lutz, Florence Cavé-Riant, Florence C. Guibal, Marie A. Breau, Yolande Di Gioia, Pierre Olivier Couraud, Yvon E. Cayre, Sandrine Bourdoulous and Pierre G. Lutz

Updated information and services can be found at:

<http://www.bloodjournal.org/content/102/9/3371.full.html>

Articles on similar topics can be found in the following Blood collections

[Cell Adhesion and Motility](#) (790 articles)

[Gene Expression](#) (1086 articles)

[Neoplasia](#) (4182 articles)

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>

Blood (print ISSN 0006-4971, online ISSN 1528-0020), is published weekly by the American Society of Hematology, 2021 L St, NW, Suite 900, Washington DC 20036.

Copyright © 2003 by The American Society of Hematology by The American Society of Hematology; all rights reserved.