

Perspective

The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes

James W. Vardiman,¹ Jüergen Thiele,² Daniel A. Arber,³ Richard D. Brunning,⁴ Michael J. Borowitz,⁵ Anna Porwit,⁶ Nancy Lee Harris,⁷ Michelle M. Le Beau,⁸ Eva Hellström-Lindberg,⁹ Ayalew Tefferi,¹⁰ and Clara D. Bloomfield¹¹

¹Department of Pathology, University of Chicago, IL; ²Institute of Pathology, University of Cologne, Cologne, Germany; ³Department of Pathology, Stanford University Medical Center, CA; ⁴Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis; ⁵Department of Pathology and Oncology, Johns Hopkins Medical Institutions, Baltimore, MD; ⁶Department of Pathology, Karolinska University Hospital, Stockholm, Sweden; ⁷Department of Pathology, Massachusetts General Hospital and Harvard Medical School, Boston; ⁸Section of Hematology/Oncology, University of Chicago, IL; ⁹Department of Medicine, Karolinska University Hospital, Stockholm, Sweden; ¹⁰Division of Hematology, Mayo Clinic, Rochester, MN; and ¹¹Comprehensive Cancer Center, The Ohio State University, Columbus

Recently the World Health Organization (WHO), in collaboration with the European Association for Haematopathology and the Society for Hematopathology, published a revised and updated edition of the *WHO Classification of Tumors of the Hematopoietic and Lymphoid Tissues*. The 4th edition of the WHO classification incorporates new information that

has emerged from scientific and clinical studies in the interval since the publication of the 3rd edition in 2001, and includes new criteria for the recognition of some previously described neoplasms as well as clarification and refinement of the defining criteria for others. It also adds entities—some defined principally by genetic features—that have only recently

been characterized. In this paper, the classification of myeloid neoplasms and acute leukemia is highlighted with the aim of familiarizing hematologists, clinical scientists, and hematopathologists not only with the major changes in the classification but also with the rationale for those changes. (*Blood*. 2009;114:937-951)

Introduction

In 2001, the World Health Organization (WHO), in collaboration with the Society for Hematopathology and the European Association of Haematopathology, published a Classification of Tumors of the Hematopoietic and Lymphoid Tissues as part of the 3rd edition of the series, *WHO Classification of Tumors*.¹ That classification reflected a paradigm shift from previous schemes in that, for the first time, genetic information was incorporated with morphologic, cytochemical, immunophenotypic, and clinical information into diagnostic algorithms for the myeloid neoplasms. The 2001 WHO classification was prefaced with a comment predicting that future revisions would be necessary because of rapidly emerging genetic and biologic information. Recently, a revised classification has been published as part of the 4th edition of the WHO monograph series.² The aim of the revision was to incorporate new scientific and clinical information to refine diagnostic criteria for previously described neoplasms and to introduce newly recognized disease entities. Our purpose in this communication is to highlight major changes in the revised WHO classification of myeloid neoplasms and acute leukemia and to provide the rationale for those changes.

Background of the WHO classification/revision

The principles of the WHO classification have been described previously.^{3,4} Briefly, the classification uses all available information—morphology, cytochemistry, immunophenotype, genetics,

and clinical features—to define clinically significant disease entities. It is a consensus classification in which a number of experts have agreed on the classification and the diagnostic criteria used for defining the entities that compose it. Nearly 30 clinicians and clinical scientists from around the world who are recognized for their expertise in myeloid neoplasms and acute leukemia were invited to be members of the Myeloid and Acute Leukemia Clinical Advisory Committee (CAC). The CAC met with the pathology committee and made a number of recommendations that were incorporated in the revision to ensure that the classification would be clinically useful. Representatives of international consensus groups such as the International Working Group for Myelofibrosis Research and Treatment and the European Group for the Immunologic Classification of Leukemia were included in the CAC.

Proposals for revisions and for recognition of new entities for the 4th edition were based on studies published in the recent literature with the goal of providing a classification that can be used in daily clinical practice as well as serve as a common language for clinical trials and laboratory investigation. However, even during the revision process, new information was being published regarding a number of the myeloid neoplasms. To be incorporated in a universally accepted classification scheme, recent data need to “mature” and their significance needs to be widely acknowledged. Therefore, to accommodate such recent information, a number of “provisional entities” are found within the major subgroups of diseases. These are newly described or characterized disorders that are clinically and/or scientifically important and should be considered in the classification, but for which additional studies are needed to clarify their significance. It is likely that many of these

provisional categories will be incorporated as confirmed entities in the next revision process.

Guidelines for using the WHO classification of myeloid neoplasms

In the WHO classification, the term “myeloid” includes all cells belonging to the granulocytic (neutrophil, eosinophil, basophil), monocytic/macrophage, erythroid, megakaryocytic and mast cell lineages. The WHO criteria for myeloid neoplasms apply to initial diagnostic peripheral blood (PB) and bone marrow (BM) specimens obtained prior to any definitive therapy for a suspected hematologic neoplasm. Morphologic, cytochemical, and/or immunophenotypic features are used for establishing the lineage of the neoplastic cells and for assessment of their maturation. The blast percentage remains a practical tool for categorizing myeloid neoplasms and judging their progression. In the WHO scheme, a myeloid neoplasm with 20% or more blasts in the PB or BM is considered to be acute myeloid leukemia (AML) when it occurs de novo, evolution to AML when it occurs in the setting of a previously diagnosed myelodysplastic syndrome (MDS) or myelodysplastic/myeloproliferative neoplasm (MDS/MPN), or blast transformation in a previously diagnosed myeloproliferative neoplasm (MPN). In some cases associated with specific genetic abnormalities, however, the diagnosis of AML may be made regardless of the blast count in the PB or BM (see “Acute myeloid leukemia and related precursor neoplasms”). The 20% blast threshold is not a mandate to treat the patient as having AML or blast transformation; therapeutic decisions must always be based on the clinical situation after all information is considered. A tumoral proliferation of blasts in an extramedullary site (myeloid sarcoma) is also considered to be AML when it is found de novo or in a patient with MDS or MDS/MPN, and blast transformation in cases of MPN. Blast percentages should be derived, when possible, from 200-cell leukocyte differential counts of the PB smear and 500-cell differential counts of all nucleated BM cells on cellular marrow aspirate smears stained with Wright-Giemsa (see Table 1). Blasts are defined using the criteria recently proposed by the International Working Group on Morphology of Myelodysplastic Syndrome.⁵ Determination of the blast percentage by assessment of CD34⁺ cells by flow cytometry is not recommended as a substitute for visual inspection; not all leukemic blasts express CD34, and hemodilution and processing artefacts can produce misleading results. For acute leukemia, multiparameter flow cytometry (3 or more colors) is the method of choice for determining the blast lineage as well as for detecting aberrant antigenic profiles that may prove useful for disease monitoring.

Although a BM core biopsy may not be required in every case, an adequate biopsy does provide the most accurate assessment of the marrow cellularity, topography, stromal changes, and maturation pattern of the hematopoietic lineages, and it can be invaluable in detecting residual disease following therapy. In addition, the biopsy provides material for immunohistochemical detection of antigens that can be diagnostically and prognostically useful, such as CD34, TdT, and Ki67, particularly if marrow aspirate smears are poorly cellular.⁶

A complete cytogenetic analysis of BM cells is essential during initial evaluation for establishing a baseline karyotype; repeat analyses are recommended as needed thereafter for judging the response to therapy or for detecting genetic evolution. Additional genetic studies should be guided by the results of the initial

Table 1. Guidelines for using the revised WHO classification of myeloid neoplasms

Specimen requirements

- PB and BM specimens collected prior to any definitive therapy.
- PB and cellular BM aspirate smears and/or touch preparations stained with Wright-Giemsa or similar stain.
- BM biopsy, at least 1.5 cm in length and at right angles to the cortical bone, is recommended for all cases if feasible.
- BM specimens for complete cytogenetic analysis and, when indicated, for flow cytometry, with an additional specimen cryopreserved for molecular genetic studies. The latter studies should be performed based on initial karyotypic, clinical, morphologic, and immunophenotypic findings.

Assessment of blasts

- Blast percentage in PB and BM is determined by visual inspection.
- Myeloblasts, monoblasts, promonocytes, megakaryoblasts (but not dysplastic megakaryocytes) are counted as blasts when summing blast percentage for diagnosis of AML or blast transformation; count abnormal promyelocytes as “blast equivalents” in APL.
- Proerythroblasts are not counted as blasts except in rare instances of “pure” acute erythroleukemia.
- Flow cytometric assessment of CD34⁺ cells is not recommended as a substitute for visual inspection; not all blasts express CD34, and artifacts introduced by specimen processing may result in erroneous estimates.
- If the aspirate is poor and/or marrow fibrosis is present, IHC on biopsy sections for CD34 may be informative if blasts are CD34⁺.

Assessment of blast lineage

- Multiparameter flow cytometry (at least 3 colors) is recommended; panel should be sufficient to determine lineage as well as aberrant antigen profile of neoplastic population.
- Cytochemistry, such as myeloperoxidase or nonspecific esterase, may be helpful, particularly in AML, NOS, but it is not essential in all cases.
- IHC on biopsy may be helpful; many antibodies are now available for recognition of myeloid and lymphoid antigens.

Assessment of genetic features

- Complete cytogenetic analysis from BM at initial diagnosis when possible.
- Additional studies, such as FISH, RT-PCR, mutational status, should be guided by clinical, laboratory, and morphologic information.
- Mutational studies for mutated *NPM1*, *CEBPA*, and *FLT3* are recommended in all cytogenetically normal AML; mutated *JAK2* should be sought in *BCR-ABL1*-negative MPN, and mutational analysis for *KIT*, *NRAS*, *PTPN11*, etc, should be performed as clinically indicated.

Correlation/reporting of data

- All data should be assimilated into one report that states the WHO diagnosis.

WHO indicates World Health Organization; PB, peripheral blood; BM, bone marrow; IHC, immunohistochemistry; AML, acute myeloid leukemia; APL, acute promyelocytic leukemia; NOS, not otherwise specified; FISH, fluorescence in situ hybridization; RT-PCR, reverse transcriptase–polymerase chain reaction; and MPN, myeloproliferative neoplasm.

karyotype and by the diagnosis suspected based on the clinical, morphologic, and immunophenotypic studies. In some cases, reverse transcriptase–polymerase chain reaction (RT-PCR) and/or fluorescence in situ hybridization (FISH) may detect variants of well-recognized cytogenetic abnormalities or submicroscopic abnormalities not detected by routine karyotyping, such as the *FIP1L1-PDGFR* fusion in some myeloid neoplasms associated with eosinophilia.⁷ In addition, gene mutations are increasingly being recognized as important diagnostic and prognostic markers in myeloid neoplasms. These include, among others, mutations of *JAK2*, *MPL*, and *KIT* in MPN⁸⁻¹³; *NRAS*, *KRAS*, *NF1*, and *PTPN11* in MDS/MPN¹⁴⁻¹⁹; *NPM1*, *CEBPA*, *FLT3*, *RUNX1*, *KIT*, *WT1*, and *MLL* in AML²⁰⁻²⁴; and *GATA1*²⁵ in myeloid proliferations associated with Down syndrome. Many of these gene mutations figure importantly in the revised WHO classification. Although over- and underexpression of genes has proved to affect the prognosis in some myeloid neoplasms,²¹ at the present time analyses of gene

Table 2. WHO classification of myeloid neoplasms and acute leukemia

Myeloproliferative neoplasms (MPN)
Chronic myelogenous leukemia, <i>BCR-ABL1</i> -positive
Chronic neutrophilic leukemia
Polycythemia vera
Primary myelofibrosis
Essential thrombocythemia
Chronic eosinophilic leukemia, not otherwise specified
Mastocytosis
Myeloproliferative neoplasms, unclassifiable
Myeloid and lymphoid neoplasms associated with eosinophilia and abnormalities of <i>PDGFRA</i>, <i>PDGFRB</i>, or <i>FGFR1</i>
Myeloid and lymphoid neoplasms associated with <i>PDGFRA</i> rearrangement
Myeloid neoplasms associated with <i>PDGFRB</i> rearrangement
Myeloid and lymphoid neoplasms associated with <i>FGFR1</i> abnormalities
Myelodysplastic/myeloproliferative neoplasms (MDS/MPN)
Chronic myelomonocytic leukemia
Atypical chronic myeloid leukemia, <i>BCR-ABL1</i> -negative
Juvenile myelomonocytic leukemia
Myelodysplastic/myeloproliferative neoplasm, unclassifiable
<i>Provisional entity: refractory anemia with ring sideroblasts and thrombocytosis</i>
Myelodysplastic syndrome (MDS)
Refractory cytopenia with unilineage dysplasia
Refractory anemia
Refractory neutropenia
Refractory thrombocytopenia
Refractory anemia with ring sideroblasts
Refractory cytopenia with multilineage dysplasia
Refractory anemia with excess blasts
Myelodysplastic syndrome with isolated del(5q)
Myelodysplastic syndrome, unclassifiable
Childhood myelodysplastic syndrome
<i>Provisional entity: refractory cytopenia of childhood</i>
Acute myeloid leukemia and related neoplasms
Acute myeloid leukemia with recurrent genetic abnormalities
AML with t(8;21)(q22;q22); <i>RUNX1-RUNX1T1</i>
AML with inv(16)(p13.1q22) or t(16;16)(p13.1;q22); <i>CBFB-MYH11</i>
APL with t(15;17)(q22;q12); <i>PML-RARA</i>
AML with t(9;11)(p22;q23); <i>MLL3-MLL</i>
AML with t(6;9)(p23;q34); <i>DEK-NUP214</i>
AML with inv(3)(q21q26.2) or t(3;3)(q21;q26.2); <i>RPN1-EV11</i>
AML (megakaryoblastic) with t(1;22)(p13;q13); <i>RBM15-MKL1</i>
<i>Provisional entity: AML with mutated NPM1</i>
<i>Provisional entity: AML with mutated CEBPA</i>
Acute myeloid leukemia with myelodysplasia-related changes
Therapy-related myeloid neoplasms
Acute myeloid leukemia, not otherwise specified
AML with minimal differentiation
AML without maturation
AML with maturation
Acute myelomonocytic leukemia
Acute monoblastic/monocytic leukemia
Acute erythroid leukemia
Pure erythroid leukemia
Erythroleukemia, erythroid/myeloid
Acute megakaryoblastic leukemia
Acute basophilic leukemia
Acute panmyelosis with myelofibrosis
Myeloid sarcoma
Myeloid proliferations related to Down syndrome
Transient abnormal myelopoiesis
Myeloid leukemia associated with Down syndrome
Blastic plasmacytoid dendritic cell neoplasm

Table 2. WHO classification of myeloid neoplasms and acute leukemia (continued)

Acute leukemias of ambiguous lineage
Acute undifferentiated leukemia
Mixed phenotype acute leukemia with t(9;22)(q34;q11.2); <i>BCR-ABL1</i>
Mixed phenotype acute leukemia with t(v;11q23); <i>MLL</i> rearranged
Mixed phenotype acute leukemia, B-myeloid, NOS
Mixed phenotype acute leukemia, T-myeloid, NOS
<i>Provisional entity: natural killer (NK) cell lymphoblastic leukemia/lymphoma</i>
B lymphoblastic leukemia/lymphoma
B lymphoblastic leukemia/lymphoma, NOS
B lymphoblastic leukemia/lymphoma with recurrent genetic abnormalities
B lymphoblastic leukemia/lymphoma with t(9;22)(q34;q11.2); <i>BCR-ABL1</i>
B lymphoblastic leukemia/lymphoma with t(v;11q23); <i>MLL</i> rearranged
B lymphoblastic leukemia/lymphoma with t(12;21)(p13;q22) <i>TEL-AML1 (ETV6-RUNX1)</i>
B lymphoblastic leukemia/lymphoma with hyperdiploidy
B lymphoblastic leukemia/lymphoma with hypodiploidy
B lymphoblastic leukemia/lymphoma with t(5;14)(q31;q32) <i>IL3-IGH</i>
B lymphoblastic leukemia/lymphoma with t(1;19)(q23;p13.3); <i>TCF3-PBX1</i>
T lymphoblastic leukemia/lymphoma

dosage by quantitative RT-PCR is not practical on a daily basis, nor have gene expression arrays been introduced into routine use, and therefore such data are not included in this revision.

Revised WHO classification of myeloid neoplasms and acute leukemia

Table 2 lists the major subgroups of myeloid neoplasms and acute leukemia in the WHO classification, and the specific entities of which they are composed. The nomenclature for the myeloproliferative entities is changed from “chronic myeloproliferative disease” to “myeloproliferative neoplasms” in order to accurately reflect their neoplastic nature. To be consistent, the subgroup formerly designated as “myelodysplastic/myeloproliferative diseases” has been renamed “myelodysplastic/myeloproliferative neoplasms.” In addition, a new subgroup, “Myeloid and lymphoid neoplasms with eosinophilia and abnormalities of *PDGFRA*, *PDGFRB*, and *FGFR1*” has been added. Within each of the major subgroups of myeloid neoplasms and acute leukemia, new entities have been added and/or diagnostic criteria have been changed.

Myeloproliferative neoplasms and myeloid neoplasms associated with eosinophilia

The MPNs are listed in Table 2. The revisions in the criteria for the WHO classification of MPN have been influenced by 2 factors: (1) the discovery of genetic abnormalities that can be used as diagnostic markers in *BCR-ABL1*-negative MPN and (2) better characterization of histologic features that aid in the identification of MPN subtypes.

In the previous WHO scheme, detection of the Philadelphia (Ph) chromosome and/or *BCR-ABL1* fusion gene was used to confirm the diagnosis of CML, whereas the *BCR-ABL1*-negative MPN subtypes were diagnosed according to their clinical and laboratory features supported by minor contributions from histopathology.¹ A number of criteria were required for distinguishing the MPN subtypes not only from each other, but also from reactive granulocytic, erythroid, and/or megakaryocytic hyperplasia that often mimics MPN. Now it is recognized that in the other MPNs, as

Table 3. Criteria for polycythemia vera (PV)

Diagnosis requires the presence of both major criteria and one minor criterion or the presence of the first major criterion together with two minor criteria:

Major criteria

1. Hemoglobin > 18.5 g/dL in men, 16.5 g/dL in women or other evidence of increased red cell volume*
2. Presence of *JAK2* V617F or other functionally similar mutation such as *JAK2* exon 12 mutation

Minor criteria

1. Bone marrow biopsy showing hypercellularity for age with trilineage growth (panmyelosis) with prominent erythroid, granulocytic, and megakaryocytic proliferation
2. Serum erythropoietin level below the reference range for normal
3. Endogenous erythroid colony formation in vitro

*Hemoglobin or hematocrit > 99th percentile of method-specific reference range for age, sex, altitude of residence
or hemoglobin > 17 g/dL in men, 15 g/dL in women if associated with a documented and sustained increase of at least 2 g/dL from a person's baseline value that cannot be attributed to correction of iron deficiency
or elevated red cell mass > 25% above mean normal predicted value.

in CML, the abnormal proliferation is due to clonal rearrangements or mutations of genes that encode surface or cytoplasmic protein tyrosine kinases (PTKs) that lead to constitutively activated signal transduction pathways.^{8-13,26,27} In some cases, the genetic abnormality, such as the *BCR-ABL1* fusion gene in CML, is associated with consistent clinical, laboratory, and/or morphologic features that allow the genetic abnormality to be used as a major criterion for diagnosis. Other abnormalities, such as mutated *JAK2* or *KIT*, are not specific for any single MPN but provide proof that the proliferation is clonal and thus, when present, eliminate further consideration of a reactive process.

At the present time, the most commonly recognized mutation in *BCR-ABL1*-negative MPN is *JAK2* V617F.^{11,26-28} This mutation is found in more than 90% of patients with polycythemia vera (PV) and in nearly one-half of those with primary myelofibrosis (PMF)

Table 4. Criteria for essential thrombocythemia (ET)

Diagnosis requires meeting all 4 criteria

1. Sustained platelet count $\geq 450 \times 10^9/L^*$
2. Bone marrow biopsy specimen showing proliferation mainly of the megakaryocytic lineage with increased numbers of enlarged, mature megakaryocytes. No significant increase or left-shift of neutrophil granulopoiesis or erythropoiesis.
3. Not meeting WHO criteria for polycythemia vera,[†] primary myelofibrosis,[‡] *BCR-ABL1*-positive CML,[§] or myelodysplastic syndrome,^{||} or other myeloid neoplasm.
4. Demonstration of *JAK2* V617F or other clonal marker, or in the absence of *JAK2* V617F, no evidence of reactive thrombocytosis[¶].

ET indicates essential thrombocythemia; BM, bone marrow; WHO, World Health Organization; and CML, chronic myelogenous leukemia.

*Sustained during the work-up process.

[†]Requires the failure of iron replacement therapy to increase hemoglobin level to the polycythemia vera range in the presence of decreased serum ferritin. Exclusion of polycythemia vera is based on hemoglobin and hematocrit levels, and red cell mass measurement is not required.

[‡]Requires the absence of relevant reticulin fibrosis, collagen fibrosis, peripheral blood leukoerythroblastosis, or markedly hypercellular marrow accompanied by megakaryocyte morphology that is typical for primary myelofibrosis—small to large megakaryocytes with an aberrant nuclear/cytoplasmic ratio and hyperchromatic, bulbous, or irregularly folded nuclei and dense clustering.

[§]Requires the absence of *BCR-ABL1*.

^{||}Requires the absence of dyserythropoiesis and dysgranulopoiesis.

[¶]Causes of reactive thrombocytosis include iron deficiency, splenectomy, surgery, infection, inflammation, connective tissue disease, metastatic cancer, and lymphoproliferative disorders. However, the presence of a condition associated with reactive thrombocytosis does not exclude the possibility of ET if other criteria are met.

Table 5. Criteria for primary myelofibrosis (PMF)

Diagnosis requires meeting all 3 major criteria and 2 minor criteria

Major criteria

1. Presence of megakaryocyte proliferation and atypia,* usually accompanied by either reticulin or collagen fibrosis,
or,
in the absence of significant reticulin fibrosis, the megakaryocyte changes must be accompanied by an increased bone marrow cellularity characterized by granulocytic proliferation and often decreased erythropoiesis (ie, prefibrotic cellular-phase disease)
2. Not meeting WHO criteria for polycythemia vera,[†] *BCR-ABL1*-positive chronic myelogenous leukemia,[‡] myelodysplastic syndrome,[§] or other myeloid disorders
3. Demonstration of *JAK2* V617F or other clonal marker (eg, *MPLW515K/L*),
or,
in the absence of the above clonal markers, no evidence that bone marrow fibrosis is secondary to infection, autoimmune disorder or other chronic inflammatory condition, hairy cell leukemia or other lymphoid neoplasm, metastatic malignancy, or toxic (chronic) myelopathies^{||}

Minor criteria

1. Leukoerythroblastosis[¶]
2. Increase in serum lactate dehydrogenase level[¶]
3. Anemia[¶]
4. Palpable splenomegaly[¶]

*Small to large megakaryocytes with an aberrant nuclear/cytoplasmic ratio and hyperchromatic, bulbous, or irregularly folded nuclei and dense clustering.

[†]Requires the failure of iron replacement therapy to increase hemoglobin level to the polycythemia vera range in the presence of decreased serum ferritin. Exclusion of polycythemia vera is based on hemoglobin and hematocrit levels. Red cell mass measurement is not required.

[‡]Requires the absence of *BCR-ABL1*.

[§]Requires the absence of dyserythropoiesis and dysgranulopoiesis.

^{||}It should be noted that patients with conditions associated with reactive myelofibrosis are not immune to primary myelofibrosis, and the diagnosis should be considered in such cases if other criteria are met.

[¶]Degree of abnormality could be borderline or marked.

or essential thrombocythemia (ET).^{11,27} Therefore, *JAK2* V617F is not specific for any single MPN, nor does its absence exclude any MPN. In the few PV patients who lack this mutation, a similar activating *JAK2* exon 12 mutation may be found,²⁹ and a small proportion of patients with PMF and ET who lack mutated *JAK2* may instead demonstrate activating mutations of *MPL*, such as *MPL* W515K or *MPL* W515L.¹² The WHO diagnostic algorithms for PV, ET, and PMF require analysis for mutated *JAK2* and if absent, for other relevant genetic abnormalities that may provide proof of clonality of the proliferative process. Still, additional parameters are required to ensure accurate diagnosis and subclassification regardless of whether a mutation is or is not found. Clinical and laboratory findings form the backbone of these diagnostic criteria, but histopathologic findings characteristic of each MPN subtype have been included in the criteria as well.³⁰ The revised criteria for PV, ET, and PMF are listed in Tables 3, 4 and 5, respectively. For ET some investigators have argued that patients with clinical and morphologic findings of ET may have platelet counts that exceed the normal range, but that fail to meet the diagnostic threshold previously required.³¹⁻³³ In the revised criteria, the platelet threshold for the diagnosis of ET has therefore been lowered from $600 \times 10^9/L$ to $450 \times 10^9/L$.

Recently, some authorities have raised concerns regarding the inclusion of histologic features as diagnostic parameters for PV, ET, and PMF. In particular, the histologic distinction between the “prefibrotic” stage of PMF (a stage often associated with marked thrombocytosis in the PB and granulocytic and atypical megakaryocytic proliferation with minimal, if any, fibrosis in the BM) and ET

has been questioned.³⁴ It is important to note that the histopathologic features included in the WHO criteria for PV, PMF, and ET were derived from a number of clinical-pathologic studies³⁵⁻³⁷ and that the WHO classification does not advocate that any single histologic feature is sufficient for a diagnosis of any MPN, but must be interpreted in the context of other clinical and laboratory criteria.

Since the last edition of the WHO classification, it has been appreciated that some cases of eosinophilia, including cases formerly designated as chronic eosinophilic leukemia (CEL) or the “hypereosinophilic syndrome,” are caused by abnormalities in genes that encode the alpha or beta moieties of the platelet-derived growth factor receptor (PDGFR). Rearrangements of *PDGFRB* at chromosome band 5q33 were first recognized in cases variably reported as chronic myelomonocytic leukemia (CMML) with eosinophilia, or less commonly, as CEL.³⁸⁻⁴¹ More recently, the gene that encodes the alpha chain of PDGFR, *PDGFRA*, located at chromosome band 4q12, was found to be involved in cryptic translocations in cases of CEL and in a substantial number of cases reported as idiopathic hypereosinophilic syndrome.⁷ Rearrangements of *FGFR1* have also been implicated in myeloproliferations with prominent eosinophilia, that is, the “8p11.2 myeloproliferative syndrome.”⁴² Patients with *FGFR1* rearrangements, however, may initially have T- or B-lymphoblastic leukemia/lymphoma associated with prominent tissue eosinophilia that may later evolve to a myeloid neoplasm with eosinophilia.^{42,43} Rare cases associated with *PDGFRA* rearrangements have also been reported to initially have a lymphoblastic neoplasm.⁴⁴ Therefore, although it might seem less confusing to categorize cases with rearranged *PDGFRA*, *PDGFRB*, or *FGFR1* and eosinophilia as CEL within the MPN category, this would ignore cases with *PDGFRB* rearrangements that have CMML as well as cases with rearrangements of *FGFR1* and *PDGFRA* that may have a lymphoid component. For these reasons, as well as for the therapeutic implications of specific identification of rearranged *PDGFRA* and *PDGFRB* that are sensitive to imatinib therapy, these cases were assigned to a new subgroup, “Myeloid/lymphoid neoplasms with eosinophilia associated with rearrangements of *PDGFRA*, *PDGFRB*, or *FGFR1*.” Cases of myeloid neoplasms with eosinophilia that lack these rearrangements should be classified as CEL, not otherwise specified (NOS), if the following criteria are met: eosinophil count is $1.5 \times 10^9/L$ or greater, blasts are less than 20% in the PB and BM, there is no *BCR-ABL1* fusion gene, *inv(16)(p13.1q22)* or *t(16;16)(p13.1;q22)*, and no evidence of another MPN or MDS/MPN, but there is a clonal, myeloid-related cytogenetic or molecular genetic abnormality, or blast cells are more than 2% in the PB or 5% in the BM. Cases with eosinophilia that lack evidence of clonality may be diagnosed as “idiopathic hypereosinophilic syndrome” after all causes of reactive eosinophilia have been excluded.⁴⁰

The appreciation of the role abnormal PTKs play in the pathogenesis of the MPNs argued for the inclusion of other chronic myeloid proliferations related to constitutively activated PTKs under the MPN umbrella. Thus, systemic mastocytosis, which has many myeloproliferative features and is almost always associated with the *KIT* D816V mutation,⁴⁵ has been added to the MPN category. Still, the pathogenesis of nearly one-half of cases of ET and PMF, all cases of chronic neutrophilic leukemia, and a number of cases of CEL remains unknown. For these, reliance on clinical, morphologic, and laboratory features is essential for diagnosis and classification.

Significant changes in the diagnosis and classification of MPN

1. The nomenclature has been changed from “myeloproliferative disorders” (MPDs) to “myeloproliferative neoplasms” (MPNs).
2. Diagnostic algorithms for PV, ET, and PMF have substantially changed to include information regarding *JAK2* V617F and similar activating mutations. Additional clinical, laboratory, and histologic parameters have been included to allow diagnosis and subclassification regardless of whether *JAK2* V617F or a similar mutation is or is not present.
3. The platelet count threshold for the diagnosis of ET has been lowered from $\geq 600 \times 10^9/L$ to $\geq 450 \times 10^9/L$.
4. Some cases previously meeting the criteria for chronic eosinophilic leukemia (CEL) may be categorized in a new subgroup, “Myeloid and lymphoid neoplasms with eosinophilia and abnormalities of *PDGFRA*, *PDGFRB*, or *FGFR1*.”
5. Systemic mastocytosis has been included in the MPN category.

Myelodysplastic/myeloproliferative neoplasms

The MDS/MPN category was introduced in the 3rd edition to include myeloid neoplasms with clinical, laboratory, and morphologic features that overlap MDS and MPN. This subgroup includes chronic myelomonocytic leukemia (CMML), atypical chronic myeloid leukemia (aCML), juvenile myelomonocytic leukemia (JMML), and a provisional entity within the MDS/MPN unclassifiable group, refractory anemia with ring sideroblasts and thrombocytosis (RARS-T). A few cases of CMML and aCML have been reported to demonstrate *JAK2* mutations,^{27,46,47} but the proliferative aspects of most cases are related to aberrancies in the RAS/MAPK signaling pathways. In JMML, nearly 75% of patients demonstrate mutually exclusive mutations of *PTPN11*, *NRAS* or *KRAS*, or *NFI*, all of which encode signaling proteins in RAS dependent pathways.^{14,16} Approximately 30% to 40% of cases of CMML and aCML also exhibit *NRAS* or *KRAS* mutations.^{17-19,48} There is therefore currently no convincing evidence to suggest that the MDS/MPN entities should be reclassified as either MPN or MDS, and the “mixed” MDS/MPN subgroup remains in the revised classification (Table 2). The distinction between CMML-1 and CMML-2, based on the percentage of blasts plus promonocytes in the PB and BM (CMML1 = < 5% blasts plus promonocytes in the PB and < 10% blasts plus promonocytes in the BM; CMML2 = 5% or more PB blasts plus promonocytes or 10% or more BM blasts plus promonocytes) suggested in the 3rd edition of the WHO classification has proved to be clinically significant^{49,50} and is again recommended in the revised edition. Cases of CMML with eosinophilia should be investigated for the *PDGFRB* abnormality and if it is found, the case should be classified as a myeloid neoplasm with eosinophilia associated with *PDGFRB* rearrangement.

During the revision process, the greatest controversy regarding MDS/MPN was related to the provisional entity RARS-T. A substantial proportion of cases of RARS-T are reported to demonstrate *JAK2* V617F or, much less commonly, mutated *MPL* (reviewed in Hellström-Lindberg⁵¹), and the percentage of patients with RARS-T who have such mutations have been reported to vary directly with the height of the platelet count.⁵² These findings have prompted some authorities to suggest that RARS-T should be moved to the MPN category, whereas others argued that RARS-T is not a distinct entity at all, but rather an MPN that has undergone genetic

evolution to acquire ring sideroblasts. The few cases of RARS-T that have been studied by in vitro culture, however, including some with *JAK2* V617F, have shown poor colony formation with lack of endogenous growth—features that are more in keeping with MDS than with MPN.^{53,54} Taken together, the *JAK2* data and the in vitro culture studies suggest that RARS-T is probably a distinct disease that is appropriately classified in the MDS/MPN category. If so, the major question regarding RARS-T involves the precise criteria for its definition, and in particular, how to distinguish it from RARS, in which modestly elevated platelet counts are often reported. This point is even more important in view of the revised criteria for RARS-T that lower the platelet threshold from $600 \times 10^9/L$ to $450 \times 10^9/L$ (in keeping with the revised criteria for ET). The revised criteria for RARS-T require not only that there be refractory anemia accompanied by thrombocytosis ($\geq 450 \times 10^9/L$) and dyserythropoiesis including ring sideroblasts that account for 15% or more of the erythroid precursors, but also that there is also a proliferation of large megakaryocytes, resembling those in ET or PMF. Such megakaryocytes are not likely to be seen in RARS. Still, because more studies are necessary for a clear definition of the boundaries of this disease, particularly the lower platelet threshold, RARS-T remains as a provisional entity in the revised classification.

Significant changes in the diagnosis and classification of MDS/MPN

1. Some cases of CMML with eosinophilia are relocated to the category “Myeloid/lymphoid neoplasms with eosinophilia and *PDGFRB* rearrangement.”
2. The category atypical CML has been renamed atypical CML, *BCR-ABL1*-negative to emphasize that it is not merely a variant of CML, *BCR-ABL1*-positive.
3. RARS-T remains as a “provisional entity” classified as MDS/MPN, unclassifiable, RARS-T. The criteria have been modified to include refractory anemia, dyserythropoiesis in the bone marrow with ring sideroblasts accounting for 15% or more of erythroid precursors, and megakaryocytes with features resembling those in PMF or ET; the platelet threshold is lowered to $450 \times 10^9/L$.

Myelodysplastic syndromes

Myelodysplastic syndromes (MDS) remain among the most challenging of the myeloid neoplasms to diagnose and classify, particularly in cases in which the blast percentage is not increased in the PB or BM. Diagnostic problems can arise when the clinical and laboratory findings suggest MDS but the morphologic findings are inconclusive; when there is secondary dysplasia caused by nutritional deficiencies, medications, toxins, growth factor therapy, inflammation or infection; or when marrow hypocellularity or myelofibrosis obscures the underlying disease process.^{6,55-58} In addition, for occasional patients in whom the diagnosis of MDS is clear, there may be difficulties in subclassification according to the previous WHO scheme, for example, in patients with isolated refractory thrombocytopenia accompanied by unilineage dysplasia of megakaryocytes. Although some diagnostic issues can be addressed in individual patients only by clinical evaluation and follow-up, the members of the CAC recommended that the minimal diagnostic criteria be more clearly stated, particularly for cases in which there is no increase in the blast percentage in the PB and/or BM, that the diagnostic role of cytogenetics and flow cytometry be clarified, and that the classification be modified to

Table 6. Recurring chromosomal abnormalities considered as presumptive evidence of MDS in the setting of persistent cytopenia of undetermined origin, but in the absence of definitive morphologic features of MDS

Unbalanced abnormalities	Balanced abnormalities
−7 or del(7q)	t(11;16)(q23;p13.3)
−5 or del(5q)	t(3;21)(q26.2;q22.1)
i(17q) or t(17p)	t(1;3)(p36.3;q21.1)
−13 or del(13q)	t(2;11)(p21;q23)
del(11q)	inv(3)(q21q26.2)
del(12p) or t(12p)	t(6;9)(p23;q34)
del(9q)	
idic(X)(q13)	

Complex karyotype (3 or more chromosomal abnormalities) involving one or more of the above abnormalities.

allow for more clinically relevant classification of patients who currently may fall into the “MDS, unclassifiable” category. Lastly, the CAC suggested that the WHO monograph should address diagnostic problems related to hypocellular MDS and MDS with myelofibrosis (MDS-F).

The “minimal” morphologic criteria for the diagnosis of MDS remain similar to those stated in the 3rd edition: in the appropriate clinical setting, at least 10% of the cells of at least one myeloid BM lineage (erythroid, granulocytic, megakaryocytic) must show unequivocal dysplasia for the lineage to be considered as dysplastic.^{56,59,60} Causes of secondary dysplasia as well as congenital abnormalities such as congenital dyserythropoietic anemia should be excluded before a diagnosis of MDS is rendered. If, however, a patient with clinical and other laboratory features consistent with MDS has inconclusive morphologic features, a presumptive diagnosis of MDS can be made if a specific clonal chromosomal abnormality, listed in Table 6, is present. It is important to note that some recurring cytogenetic abnormalities observed in MDS, particularly del(20q), +8, and −Y, are not included in the list. These abnormalities have been reported to occur in some patients with aplastic anemia or other cytopenic syndromes who have a good response to immunosuppressive therapy and/or who show no morphologic evidence of MDS with prolonged follow-up.⁶¹⁻⁶³ Moreover, loss of the Y chromosome in hematopoietic cells has also been reported to be a phenomenon associated with aging.⁶⁴ Thus, it is not yet clear that these abnormalities are necessarily indicative of MDS when the morphologic features are not conclusive. If only unilineage dysplasia is present in the BM but one of the recurrent cytogenetic abnormalities listed in Table 6 is not present, and there is no increase in blasts in the PB or BM, and ring sideroblasts are less than 15% of the erythroid precursors, an observation period of 6 months and repeat bone marrow investigation is recommended prior to making the diagnosis of MDS. This will ensure that the clinical and morphologic features persist and are not secondary to any other disorder that emerges in that period.

Whether phenotypic abnormalities demonstrated by flow cytometry, such as asynchronous expression of maturation-associated antigens on myeloid cells, are diagnostic of MDS when morphologic features are inconclusive was considered at the CAC meeting. The consensus was that, although numerous authors have reported aberrant antigen expression patterns characteristic for MDS cells,⁶⁵⁻⁶⁷ sufficient numbers of cases with secondary dysplasia have not been adequately evaluated so that it can be proved that the changes are specific for MDS. Until such data are available, it is recommended that, if 3 or more phenotypic abnormalities are found involving one or more of the myeloid lineages the findings can be considered as

Table 7. Peripheral blood and bone marrow findings in myelodysplastic syndrome (MDS)

Disease	Blood findings	BM findings
Refractory cytopenia with unilineage dysplasia (RCUD): (refractory anemia [RA]; refractory neutropenia [RN]; refractory thrombocytopenia [RT])	Unicytopenia or bicytopenia* No or rare blasts (< 1%)†	Unilineage dysplasia: ≥ 10% of the cells in one myeloid lineage < 5% blasts < 15% of erythroid precursors are ring sideroblasts
Refractory anemia with ring sideroblasts (RARS)	Anemia No blasts	≥ 15% of erythroid precursors are ring sideroblasts Erythroid dysplasia only < 5% blasts
Refractory cytopenia with multilineage dysplasia (RCMD)	Cytopenia(s) No or rare blasts (< 1%)† No Auer rods < 1 × 10 ⁹ /L monocytes	Dysplasia in ≥ 10% of the cells in ≥ 2 myeloid lineages (neutrophil and/or erythroid precursors and/or megakaryocytes) < 5% blasts in marrow No Auer rods ± 15% ring sideroblasts
Refractory anemia with excess blasts-1 (RAEB-1)	Cytopenia(s) < 5% blasts† No Auer rods < 1 × 10 ⁹ /L monocytes	Unilineage or multilineage dysplasia 5%-9% blasts† No Auer rods
Refractory anemia with excess blasts-2 (RAEB-2)	Cytopenia(s) 5%-19% blasts‡ Auer rods ±‡ < 1 × 10 ⁹ /L monocytes	Unilineage or multilineage dysplasia 10%-19% blasts‡ Auer rods ±‡
Myelodysplastic syndrome—unclassified (MDS-U)	Cytopenias < 1% blasts†	Unequivocal dysplasia in < 10% of cells in one or more myeloid lineages when accompanied by a cytogenetic abnormality considered as presumptive evidence for a diagnosis of MDS (see Table 6) < 5% blasts
MDS associated with isolated del(5q)	Anemia Usually normal or increased platelet count No or rare blasts (< 1%)	Normal to increased megakaryocytes with hypolobated nuclei < 5% blasts Isolated del(5q) cytogenetic abnormality No Auer rods

*Bicytopenia may occasionally be observed. Cases with pancytopenia should be classified as MDS-U.

†If the marrow myeloblast percentage is < 5% but there are 2% to 4% myeloblasts in the blood, the diagnostic classification is RAEB-1. Cases of RCUD and RCMD with 1% myeloblasts in the blood should be classified as MDS-U.

‡Cases with Auer rods and < 5% myeloblasts in the blood and less than 10% in the marrow should be classified as RAEB-2. Although the finding of 5% to 19% blasts in the blood is, in itself, diagnostic of RAEB-2, cases of RAEB-2 may have < 5% blasts in the blood if they have Auer rods or 10% to 19% blasts in the marrow or both. Similarly, cases of RAEB-2 may have < 10% blasts in the marrow but may be diagnosed by the other 2 findings, Auer rod+ and/or 5% to 19% blasts in the blood.

“suggestive” of MDS, but in the absence of conclusive morphologic and/or cytogenetic features, flow cytometric abnormalities alone are not diagnostic of MDS. Patients whose cells exhibit aberrant phenotypic features suggestive of MDS should be carefully followed for morphologic features sufficient to substantiate the diagnosis.

Despite clarifications of the diagnostic criteria for MDS, there will be a number of patients with persistent cytopenia(s) who lack sufficient morphologic or cytogenetic evidence for a definitive diagnosis of MDS. Some authors have suggested the term “idiopathic cytopenia of undetermined significance” (ICUS) for such cases.⁶⁸ Although this term is reasonable, it should be construed not as an entity in the WHO classification of MDS, but as a description for patients who do not fulfill the minimal WHO criteria. These patients should be carefully monitored for emerging morphologic or cytogenetic evidence of MDS.

The revised WHO classification of MDS has been further refined to allow for a more precise and prognostically relevant subclassification of patients with unilineage dysplasia who, in the previous scheme, were considered as “unclassifiable.” Such cases included patients with isolated thrombocytopenia with dysplasia

limited to the megakaryocytic lineage or neutropenia with only granulocytic dysplasia, and no blasts in the blood and fewer than 5% blasts in the bone marrow. The category of “refractory cytopenia with unilineage dysplasia” has been expanded and redefined to include MDS patients with less than 1% blasts in the PB and less than 5% in the BM, but 10% or more dysplastic cells in a single myeloid lineage. This category includes the entities of refractory anemia (RA) for those with anemia and only dyserythropoiesis (but in whom ring sideroblasts account for less than 15% of the erythroid precursors), refractory neutropenia (RN) for those with neutropenia and only dysgranulopoiesis, and refractory thrombocytopenia (RT) for those with thrombocytopenia and morphologic dysplasia limited to the megakaryocytic lineage. The criteria also permit patients to be included in one of these categories if the dysplasia is unilineage but there is bi-cytopenia in the PB. On the other hand, patients with pancytopenia and unilineage morphologic dysplasia are classified as having MDS, unclassifiable, because of the uncertain clinical significance of such findings. These changes are outlined in Table 7.

Some authors suggested that the previous WHO classification failed to emphasize the prognostic significance of increased

blasts in the PB.⁶⁹ To address this criticism, the revised classification redefines the criteria for refractory anemia with excess blasts (RAEB-1) to include patients with PB smears that consistently show 2% to 4% blasts in the blood, even if the blast percentage in the BM is less than 5%. Patients with 5% to 19% PB blasts or 10% to 19% blasts in the BM are classified in the highest WHO grade, RAEB-2.

A minority of patients with MDS initially have hypocellular or myelofibrotic bone marrow specimens.^{70,71} Hypoplastic MDS has no independent prognostic significance per se, but such cases can pose diagnostic problems because they may show some morphologic similarities with aplastic anemia, such as macrocytic red blood cells. Also, MDS-F can be difficult to diagnose and subclassify because cellular BM aspirate smears cannot be obtained for evaluation of either dysplasia or blast percentage. Furthermore, distinction of MDS-F from other myeloid neoplasms with fibrosis, such as acute panmyelosis with myelofibrosis and acute megakaryocytic leukemia, is often difficult. Although criteria for the diagnosis of “hypocellular MDS” and “MDS-F” have been suggested by various investigators,^{70,72} none are universally accepted nor, in the opinion of the members of the CAC and pathology committees, is it clear that they define a unique type of MDS that cannot be otherwise accommodated within the framework of current MDS criteria. Therefore, the revised WHO classification does not recognize hypocellular MDS and MDS-F as distinct entities, but recommends that, once such cases are found to meet the criteria for MDS, they should be subclassified according to the MDS guidelines, followed by “hypoplastic” or “with myelofibrosis” as modifiers to the WHO subgroup, for example, “RAEB-2 with myelofibrosis.” In MDS with hypocellularity or fibrosis, careful examination of the PB for dysplasia, estimation of CD34⁺ blasts by immunohistochemistry in the BM biopsy, and correlation with cytogenetic findings will usually allow the correct diagnosis and subclassification of the case.^{6,73,74} Most cases with myelofibrosis will demonstrate an excess of blasts that can be readily appreciated by immunohistochemical stains for CD34 on the biopsy, whereas those with hypoplastic bone marrow specimens are often in the lower grade groups. On the other hand, in a hypocellular bone marrow that shows only unilineage dysplasia—particularly of the erythroid lineage—and no cytogenetic abnormalities or increase in blasts, a diagnosis of MDS should be deferred for an observation period of 6 months if possible, as suggested for cases of RA without cytogenetic abnormalities.

The 3rd edition of the WHO classification was criticized by some pediatric hematologists for failing to recognize differences between the clinical and pathologic features of MDS in children and MDS in adults.⁷⁵ At the CAC meeting it was the consensus of the pediatric hematologists and hematopathologists that children with MDS and 2% to 19% blasts in the PB and/or 5% to 19% blasts in the BM may be categorized according to the same criteria as for adults with MDS. However, in contrast to adults, isolated refractory anemia is uncommon in childhood MDS, which more commonly is initially associated with thrombocytopenia and/or neutropenia, often accompanied by a hypocellular BM.⁷⁵ In order to emphasize the unique features of these cases of childhood MDS, the 4th edition of the WHO devotes a section to childhood MDS, in which a provisional entity, refractory cytopenia of childhood (RCC), is introduced in the classification. The category of RCC is reserved for childhood cases with less than 2% blasts in the blood and less than 5% in the marrow and persistent cytopenias associated with

dysplasia in at least 2 cell lineages. Often, BM specimens in RCC are hypocellular, and if no MDS-related cytogenetic abnormalities are present, the distinction between RCC and aplastic anemia or congenital bone marrow failure syndromes may be very difficult. Furthermore, the distinction, if any, between RCC and refractory cytopenia with multilineage dysplasia as defined in adults is not clear at this time. More study is needed to clarify these questions, thus RCC is considered as a provisional entity.

Significant changes in the diagnosis and classification of MDS

1. Patients with refractory cytopenia(s) suspected to have MDS, but who lack diagnostic morphologic features may be considered to have presumptive evidence of MDS if they have specific MDS-related cytogenetic abnormalities.
2. An over-arching category of MDS, refractory cytopenia with unilineage dysplasia, has been added to incorporate patients who exhibit unilineage dysplasia associated with refractory anemia (unilineage erythroid dysplasia), refractory neutropenia (unilineage dysgranulopoiesis), or refractory thrombocytopenia (unilineage dysmegakaryocytopoiesis), and who have fewer than 1% blasts in the blood and fewer than 5% in the bone marrow.
3. The category of refractory cytopenia with multilineage dysplasia is no longer subdivided according to whether 15% or more of the erythroid precursors are ring sideroblasts (RS), that is, the former category of RCMD-RS is now incorporated in RCMD.
4. Patients with 2% to 4% blasts in the blood and less than 5% blasts in the bone marrow should be diagnosed as having RAEB-1 if other clinical and laboratory findings of MDS are present.
5. A provisional entity, refractory cytopenia of childhood (RCC), has been added to include children with cytopenia(s) with less than 2% blasts in the peripheral blood and less than 5% in the bone marrow and evidence of dysplasia in 2 or more lineages. For children with 2% to 19% blasts in the blood and/or 5% to 19% in the bone marrow, the MDS subclassification should be made using the same criteria used for adults.

Acute myeloid leukemia and related precursor neoplasms

The previous edition of the WHO classification ushered in the era of incorporation of genetic abnormalities into diagnostic algorithms for the diagnosis of AML. The 4 abnormalities initially included in the subgroup “AML with recurring genetic abnormalities” involved rearrangements of genes (*RUNX1*, *CBFB*, *RARA*, and *MLL*) that encode transcription factors and that are associated with fairly distinct clinical and morphologic features. Since the publication of the 3rd edition of the classification, it has become more widely appreciated that multiple genetic lesions—including not only microscopically detectable chromosomal rearrangements or numerical abnormalities but also submicroscopic gene mutations—cooperate to establish the leukemic process and influence its morphologic and clinical characteristics.⁷⁶ Although rearrangements of genes that encode transcription factors may lead to impaired maturation of one or more myeloid lineages, mutations of genes such as *FLT3*, *JAK2*, *RAS*, or *KIT* that encode proteins involved in signal transduction pathways may be required for the proliferation and/or survival of the neoplastic clone. The discovery of the importance of gene mutations in leukemogenesis has also paved the way for the genetic characterization of many cases of cytogenetically normal AML.^{20,21,23,77} In cytogenetically abnormal

as well as in cytogenetically normal AML, these newly discovered genetic abnormalities may be associated with clinical, morphologic, and/or phenotypic features that allow identification of a specific leukemic entity, and in other instances have proved to be powerful prognostic indicators even when they do not define a specific leukemic subtype.⁷⁷

One of the major challenges in the revision of the WHO classification of AML was how to incorporate important and/or recently described genetic aberrations into a classification scheme of AML and yet adhere to the WHO principle of defining homogeneous, biologically relevant, and mutually exclusive entities based not only on the prognostic value of a genetic abnormality, but on morphologic, clinical, phenotypic, and/or other unique biologic properties. This was particularly problematic for the most frequent and prognostically important mutations currently recognized in cytogenetically normal AML, mutated *FLT3*, *NPM1*, and *CEBPA*. These mutations are associated with few, if any, entirely consistent morphologic or clinical features. Most notably, they are not entirely mutually exclusive of each other or, particularly in the case of *FLT3*, of other well-recognized cytogenetic defects.^{20,78} Yet, in the context of cytogenetically normal AML, they are each important prognostic factors. Mutated *NPM1* is found in 50% to 60% of cases of cytogenetically normal AML, and when only the *NPM1* mutation status is considered, is reported to be associated with approximately 50% survival at 4 years.^{21,77} The *FLT3-ITD* mutation is found in approximately 30% to 40% of cases of cytogenetically normal AML, and when only the *FLT3-ITD* status is considered is associated with a survival rate of only 20% to 25% at 4 years. Mutations of *CEBPA* are found in approximately 15% of cytogenetically normal AML cases and these patients have a nearly 60% 4-year survival rate. When *NPM1* and *FLT3-ITD* status are considered together, patients who have mutated *NPM1* and are *FLT3-ITD*-negative have a 4-year survival similar to that of the *CEBPA*-mutated cases at approximately 60%, whereas the remaining cases, being either *NPM1* wild-type or *FLT3-ITD*-positive or both, have a dismal 30% survival rate at 4 years.^{21,77}

Despite the dilemma and controversy regarding how to incorporate recently described and clinically important genetic lesions into the revised scheme, the framework of the classification proved flexible enough so that new, well-defined entities could be incorporated as could provisional entities—those for which more data need to be collected so that they can be better characterized. Table 2 shows the revised AML classification. In the subgroup of AML with recurrent genetic abnormalities, 3 entities included in the previous edition are retained: (1) AML with t(8;21)(q22;q22); *RUNX1-RUNX1T1*; (2) AML with inv(16)(p13.1;q22) or t(16;16)(p13.1;q22); *CBFB-MYH11*; and (3) acute promyelocytic leukemia (APL) with t(15;17)(q22;q12); *PML-RARA*. However, variant *RARA* translocations, such as those involving *ZBTB16* at 11q23, *NUMA1* at 11q13, *NPM1* at 5q35, or *STAT5B* at 17q11.2, should be diagnosed as AML with the variant partner specifically designated.⁷⁹ For these entities alone [AML with t(8;21)(q22;q22), AML with inv(16)(p13.1;q22) or t(16;16)(p13.1;q22), and APL with t(15;17)(q22;q12)] the genetic abnormality is sufficient for the diagnosis of AML in the appropriate setting regardless of the blast percentage in the PB or BM. For all other entities within the category of AML with recurrent genetic abnormalities, 20% or more blasts must be present in the PB or BM to establish the diagnosis of AML. Although these leukemias are recognized as unique entities, it is important to realize that additional genetic abnormalities may coexist and influence their biologic and clinical behavior, including response to therapy and the overall survival

(OS). For example, mutated *KIT* is not uncommon in core-binding factor AML, that is, those with rearranged *RUNX1* and *CBFB* genes. In the case of t(8;21)(q22;q22); *RUNX1-RUNX1T1*, most published reports indicate a higher relapse rate and lower OS when mutated *KIT* is present, whereas its impact in the presence of the inv(16)(p13.1;q22) or t(16;16)(p13.1;q22) remains to be firmly established.^{20,22} Deciding how to incorporate coexisting genetic abnormalities that are of prognostic importance or that provide a target for molecularly directed therapy will likely be increasingly important in future revisions of the classification, particularly as targeted therapies become available.

One entity included in the recurring genetic abnormality subgroup in the 3rd edition, “AML with abnormalities of 11q23; *MLL*,” has been revised to focus on the most frequent and best characterized of the *MLL* abnormalities, AML with t(9;11)(p22;q23); *MLL3-MLL*. More than 80 partner genes participate in translocations with *MLL*, and in addition, “abnormalities of *MLL*,” as used in the previous edition, would include the partial tandem duplication of the gene.⁸⁰⁻⁸² The AMLs that result from these various rearrangements and aberrations of *MLL* are not identical, and, in the case of translocations, the biology is influenced by the partner gene. Although the t(9;11)(p22;q23) is specifically listed in the classification, it is recommended that variant *MLL* translocations be also specified in the diagnosis, for example, AML with t(11;19)(q23;p13.3); *MLL-ENL*.

Studies of AML associated with the chromosomal rearrangements t(6;9)(p23;q34); *DEK-NUP21*, inv(3)(q21q26.2) or t(3;3)(q21;q26.2); *RPN1-EVII*, and t(1;22)(p13;q13); *RBM15-MKLI* have provided evidence that these genetic rearrangements, although uncommon, are associated with distinctive morphologic and clinical features that argue for their incorporation in the revised listing of AML with recurrent genetic abnormalities.⁸³⁻⁸⁵ A more controversial issue is whether leukemia initially demonstrating a blastic myeloid proliferation associated with the Ph chromosome or *BCR-ABL1* is a distinct and easily defined entity. Although *BCR-ABL1*-positive AML has been reported,⁸⁶⁻⁸⁸ criteria for its distinction from CML initially manifesting in a blast phase are not entirely convincing, and for this reason, *BCR-ABL1*-positive AML is not recognized in this classification. Many cases of *BCR-ABL1*-related acute leukemia will meet the criteria for acute lymphoblastic leukemia or mixed phenotype acute leukemia (see “Acute leukemia of ambiguous lineage”), provided that a blast phase of a previously unrecognized CML can be excluded.

In addition to the newly listed leukemias associated with cytogenetically detectable rearrangements, AML with mutated *NPM1* or *CEBPA* have been added to the classification as “provisional entities.” Because mutated *FLT3* frequently accompanies other genetic lesions, including well-recognized abnormalities such as the t(15;17)(q22;q12) and the t(6;9)(p23;q34), it is not included as a defining criterion for any distinct entity in this revision. Nevertheless, because of its prognostic importance, the mutational status of *FLT3* should always be ascertained in AML, particularly in cytogenetically normal AML.

AML with multilineage dysplasia was initially introduced in the WHO classification to encompass cases of AML characterized by MDS-like features, including unfavorable cytogenetic abnormalities, overexpression of multidrug-resistance glycoprotein, and an unfavorable response to therapy. Dysplasia in at least 50% of the cells in 2 or more hematopoietic lineages was used as a surrogate marker for these MDS-related biologic features. Although the prognostic significance of AML with multilineage dysplasia has been verified in some studies,⁸⁹⁻⁹¹ others showed that morphologic

Table 8. Cytogenetic abnormalities sufficient for diagnosis of AML with myelodysplasia-related changes when 20% or more PB or BM blasts are present

Complex karyotype*	
Unbalanced abnormalities	Balanced abnormalities
–7 or del(7q)	t(11;16)(q23;p13.3)†
–5 or	t(3;21)(q26.2;q22.1)†
i(17q) or t(17p)	t(1;3)(p36.3;q21.1)
–13 or del(13q)	t(2;11)(p21;q23)†
del(11q)	t(5;12)(q33;p12)
del(12p) or t(12p)	t(5;7)(q33;q11.2)
del(9q)	t(5;17)(q33;p13)
idic(X)(q13)	t(5;10)(q33;q21)
	t(3;5)(q25;q34)

*Three or more unrelated abnormalities, none of which are included in the “AML with recurrent genetic abnormalities” subgroup; such cases should be categorized in the appropriate cytogenetic group.

†These abnormalities most commonly occur in therapy-related disease, and therapy-related AML should be excluded before these are used as evidence for diagnosis of AML with myelodysplasia-related changes.

multilineage dysplasia had no independent prognostic significance when cytogenetic findings were incorporated in the analysis.^{92,93} To address these studies, this subgroup has been renamed as “AML with myelodysplasia-related changes,” and the criteria have expanded to include a history of MDS or MDS/MPN and cytogenetic findings. Patients may be assigned to this category if they have 20% or more blasts in the PB or BM and (1) evolve from previously documented MDS or MDS/MPN, (2) have specific myelodysplasia-related cytogenetic abnormalities (see Table 8), or (3) exhibit dysplasia in 50% or more of the cells in 2 or more myeloid lineages. The diagnosis should state the reason(s) for placing the patient in this category, for example, “AML with myelodysplasia-related changes with myelodysplasia-related cytogenetic abnormalities.” Patients who are categorized in this subgroup and have a normal karyotype should be analyzed for *FLT3*, *NPM1*, and *CEBPA* mutations, and, if present, the abnormality should be noted in the diagnosis. Currently, the significance of one or more of these mutations in the setting of morphologic dysplasia is not known.

Therapy-related myeloid neoplasms (t-AML/t-MDS and t-AML/t-MDS/MPN) remain as a distinct subgroup in the AML classification. Although cases may be designated as t-AML, or as t-MDS or t-MDS/MPN depending on the blast count, it is useful to think of them as a single biologic disease with similar genetic features, so that the designation as t-AML/t-MDS is appropriate as well.^{94,95} Most patients who develop therapy-related myeloid neoplasms have received alkylating agents and/or radiation as well as topoisomerase II inhibitors, so that a division according to the type of therapy is usually not practical and is no longer recommended. It has been argued that 90% of patients with therapy-related neoplasms have cytogenetic abnormalities identical to those observed in AML with myelodysplasia-related features or in AML with recurrent cytogenetic abnormalities, therefore such cases would be more appropriately classified in those categories rather than in a separate, therapy-related category. However, except for some patients with t-AML associated with inv(16)(p13.1q22), t(16;16)(p13.1;q22) or t(15;17)(q22;q12), patients with therapy-related myeloid neoplasms have a significantly worse outcome than do their de novo counterparts with the same genetic abnormalities, suggesting that there are biologic differences.^{94,96-99} Furthermore, the study of therapy-related neoplasms may provide valuable insight into the pathogenesis of de novo disease by providing clues as to why only a few patients develop leukemia, whereas most

patients treated with the identical agents do not. It is recommended, however, that the cytogenetic abnormality be included in the diagnostic report, for example, “therapy-related AML with t(9;11)(p22;q23).”

Acute myeloid leukemia, not otherwise specified (AML, NOS) encompasses those cases that do not fulfill the criteria for any of the other AML categories. This category accounts for only 25% to 30% of all cases, and it will continue to diminish as more genetic subgroups are recognized. One of the lively discussions at the CAC meeting was whether to recommend continued subclassification within this group by criteria similar to those of the previous French-American-British (FAB) classification scheme for AML, or to designate all such cases merely as AML, NOS. The argument that specific genetic abnormalities may yet be found that will further characterize some AML, NOS subgroups prevailed as a reason for maintaining the subclassification. Furthermore, for prognostic purposes, it may still be important to recognize some disorders such as acute megakaryoblastic leukemia, acute panmyelosis with myelofibrosis, and pure acute erythroleukemia. However, it is important to note that the information used to characterize the subgroups in this category, such as epidemiologic or clinical outcome data, is often based on older studies that included patients who would now be assigned to different WHO diagnostic categories. Therefore, such information may no longer be reliable for patients now classified in this subgroup.

Three additional myeloid neoplasms in the AML listing include myeloid sarcoma, the myeloid proliferations related to Down syndrome, and blastic plasmacytoid dendritic cell neoplasm. Myeloid sarcoma, previously known as granulocytic sarcoma or chloroma, is a pathologic diagnosis for an extramedullary proliferation of blasts of one or more of the myeloid lineages that disrupts the normal architecture of the tissue in which it is found.¹⁰⁰ Any site of the body can be affected, particularly the skin, gastrointestinal tract, lymph nodes, and bone. Most often, myeloid sarcoma is found concurrently in a patient with previously or recently recognized AML, but it may also precede the appearance of blood or bone marrow disease. In this latter case, the diagnosis of myeloid sarcoma should be considered as synonymous with AML, and the tumor should be evaluated for morphologic, phenotypic, and genetic features that would allow it to be classified further into one of the subgroups of AML. A myeloid sarcoma is evidence of relapse in a patient thought to be in remission of a previously diagnosed AML, is evidence of evolution to AML in a patient with a previously diagnosed MDS or MDS/MPN, and is blast transformation in a patient with MPN. The myeloid proliferations related to Down syndrome—transient abnormal myelopoiesis and myeloid leukemia—have unique morphologic, immunophenotypic, clinical, and molecular features, including *GATA1* mutation, that justify their separation from other myeloid neoplasms.^{25,101-103} MDS related to Down syndrome is biologically identical to Down-related AML; therefore they are considered as a single entity, myeloid leukemia associated with Down syndrome, in the classification. The neoplasm referred to in the WHO 3rd edition as “blastic NK-cell lymphoma” or in the literature as “agranular CD4⁺/CD56⁺ hematodermic neoplasm” is now known to be in virtually all cases a tumor that is derived from precursors of a specialized subset of dendritic cells, plasmacytoid dendritic cells, and hence is a myeloid-related neoplasm.¹⁰⁴⁻¹⁰⁷ It is a clinically aggressive neoplasm that is usually characterized at its onset by solitary or multiple skin lesions, often with associated regional lymphadenopathy. Many cases will ultimately progress to involve the PB and BM as well. The blasts in such cases do not express myeloperoxidase or

nonspecific esterase, and they are characterized by the expression of CD4, CD43, CD56, CD123, BDCA-2/CD303, TCL1, and CLA; CD7 and CD33 are not uncommonly expressed as well, and TdT is expressed in approximately 30% of cases. There is no expression of CD34 or CD117.^{106,107}

Significant changes in the diagnosis and classification of AML and related precursor neoplasms

1. AML with recurrent genetic abnormalities

- a. As in the previous edition, AML with t(8;21)(q22;q22), inv(16)(p13.1q22) or t(16;16)(p13.1;q22), and APL with t(15;17)(q22;q12) are considered as acute leukemia regardless of blast count in the PB or BM, but in contrast to the previous edition, for AML with t(9;11)(p22;q23) or other 11q23 abnormalities, as well as for all other subgroups (except the rare instance of some cases of erythroleukemia) blasts of 20% or more of white blood cells in PB or of all nucleated BM cells is required for the diagnosis of AML.
- b. In APL with t(15;17)(q22;q12); *PML-RARA*, variant *RARA* translocations with other partner genes are recognized separately; not all have typical APL features and some have all-*trans* retinoic acid (ATRA) resistance.
- c. The former category, AML with 11q23 (*MLL*) abnormalities has been redefined to focus on AML with t(9;11)(p22;q23); *MLLT3-MLL*. Translocations of *MLL* other than that involving *MLLT3* should be specified in the diagnosis. Other abnormalities of *MLL*, such as partial tandem duplication of *MLL* should not be placed in this category.
- d. Three new cytogenetically defined entities are added: (1) AML with t(6;9)(p23;q34); *DEK-NUP214*, (2) AML with inv(3)(q21q26.2) or t(3;3)(q21;q26.2); *RPNI-EVII*; and (3) AML (megakaryoblastic) with t(1;22)(p13;q13); *RBM15-MKLL*.
- e. Two provisional entities are added: AML with mutated *NPM1* and AML with mutated *CEBPA*. Although not included as a distinct or provisional entity, examination for mutations of *FLT3* is strongly recommended in all cases of cytogenetically normal AML.

2. AML with myelodysplasia-related changes

- a. The name was changed and expanded from “AML with multilineage dysplasia” to “AML with myelodysplasia-related changes.”
- b. Cases of AML are assigned to this category if (1) they have a history of MDS or MDS/MPN and have evolved to AML, (2) they have a myelodysplasia-related cytogenetic abnormality, or (3) at least 50% of cells in 2 or more myeloid lineages are dysplastic.

3. Therapy-related myeloid neoplasms

Cases are no longer subcategorized as “alkylating agent related” or “topoisomerase II-inhibitor related.”

4. AML, NOS

- a. Some cases previously assigned to the subcategory of AML, NOS as acute erythroid leukemia or acute megakaryoblastic leukemia may be reclassified as AML with myelodysplasia-related changes.
- b. Cases previously categorized as AML, NOS, acute megakaryoblastic leukemia should be placed in the appropriate genetic category if they are associated with inv(3)(q21q26.2) or t(3;3)(q21;q26.2); *RPNI-EVII*, or AML (megakaryoblastic) with t(1;22)(p13;q13); *RBM15-MKLL*. Down syndrome-related cases are excluded from this category as well.

Table 9. Requirements for assigning more than one lineage to a single blast population in mixed phenotype acute leukemia (MPAL)

Myeloid lineage

Myeloperoxidase (flow cytometry, immunohistochemistry, or cytochemistry)
or

Monocytic differentiation (at least 2 of the following: nonspecific esterase, CD11c, CD14, CD64, lysozyme)

T lineage

Cytoplasmic CD3 (flow cytometry with antibodies to CD3 epsilon chain; immunohistochemistry using polyclonal anti-CD3 antibody may detect CD3 zeta chain, which is not T cell-specific)
or

Surface CD3 (rare in mixed phenotype acute leukemia)

B lineage (multiple antigens required)

Strong CD19 with at least 1 of the following strongly expressed: CD79a, cytoplasmic CD22, CD10
or

Weak CD19 with at least 2 of the following strongly expressed: CD79a, cytoplasmic CD22, CD10

5. Myeloid proliferations related to Down syndrome

This new category incorporates transient abnormal myelopoiesis as well as MDS and AML that is Down syndrome-related. MDS and AML related to Down syndrome are biologically identical and thus are considered together as “Myeloid leukemia associated with Down syndrome.”

6. Blastic plasmacytic dendritic cell neoplasm

This is a new category that includes most cases previously classified as blastic NK-cell lymphoma/leukemia or agranular CD4⁺ CD56⁺ hematodermic neoplasm; it is derived from a precursor of plasmacytoid dendritic cells.

Acute leukemia of ambiguous lineage

Acute leukemias of ambiguous lineage show no clear evidence of differentiation along a single lineage. In some cases, no lineage-specific antigens are present, whereas in others the blasts express antigens of more than one lineage to such a degree that it is not possible to assign the leukemia to a specific lineage-related category. Historically, there has been confusion in the definition, terminology, and the criteria used for subclassification of such cases.¹⁰⁸⁻¹¹⁰ In an attempt to clarify the definition of this group of diseases and to simplify their diagnosis, the 4th edition of the WHO classification not only places acute leukemias of ambiguous lineage in a chapter distinct from those of AML and ALL, but has significantly altered the criteria used to define the largest subset of these cases—those that express antigens of more than one lineage. Cases with no lineage-specific markers are designated as acute undifferentiated leukemia (AUL). Such cases often express CD34, HLA-DR, and/or CD38, and sometimes TdT, but lack specific myeloid or lymphoid antigens.¹¹¹⁻¹¹³ Leukemias with blasts that coexpress certain antigens of more than one lineage on the same cells or that have separate populations of blasts that are of different lineages are referred to as mixed phenotype acute leukemia (MPAL). These cases may be further designated as B-myeloid or T-myeloid, irrespective of whether one or more than one population of blasts is found. The requirements for assigning specific lineages to the blasts are given in Table 9. Only a limited number of antigens are used in defining the pattern of lineage involvement. It should be pointed out that the requirement for myeloperoxidase positivity to prove myeloid lineage should be applied to MPAL only and not to

cases of AML with minimal differentiation that do not show any lymphoid-associated antigens. In general, leukemias that can be classified according to a specific genetic defect, such as the t(8;21)(q22;q22), are excluded from the MPAL category, with the exception of some acute leukemias associated with the *BCR-ABL1* fusion gene or with *MLL* rearrangements. In the specific case of acute leukemia with the Ph chromosome or *BCR-ABL1* fusion, if the requirements for assigning the blast population to more than one lineage are fulfilled, a diagnosis of MPAL with t(9;22)(q34;q11.2) can be made, provided the blast phase of CML can be excluded.

There is one provisional entity included in this category, “natural killer cell lymphoblastic leukemia/lymphoma.” Most cases previously designated as “blastic natural killer cell leukemia/lymphoma” are now recognized as the myeloid-related “blastic plasmacytoid dendritic cell neoplasm.”^{105,106} The defining phenotype of true NK-cell precursor neoplasms is not clear, although the diagnosis might be considered when the blasts express CD56 with immature T-associated antigens such as CD7 and CD2, in the absence of any B or myeloid antigen expression, particularly if no rearrangement of T-cell receptor genes can be detected. However, this provisional subtype needs further study and clarification.

Summary of changes in the diagnosis and classification of acute leukemia of ambiguous lineage

1. Leukemia formerly designated as “bilineal acute leukemia” and “biphenotypic acute leukemia” are now collectively considered as “mixed phenotype acute leukemia”(MPAL).
2. The criteria that define the myeloid, T-lymphoid, and B-lymphoid components of mixed phenotype acute leukemia have been significantly altered (see Table 9).
3. Cases of *BCR-ABL1*-positive and *MLL*-positive acute leukemias may meet the criteria for MPAL; in the case of *BCR-ABL1*-positive disease, CML in blast phase should be excluded.
4. Blastic natural killer cell leukemia/lymphoma is not easily defined and is considered as a provisional entity in this category; most cases previously designated as such are now recognized as blastic plasmacytoid dendritic cell neoplasms.

Precursor lymphoid neoplasms: B lymphoblastic leukemia/lymphoma and T lymphoblastic leukemia/lymphoma

The classification of precursor B-cell and T-cell neoplasms is shown in Table 2. Although the distinction between lymphoblastic leukemia and lymphoblastic lymphoma is obvious when the patient has a mass composed of B or T lymphoblasts and no blasts in the blood or marrow, it is more arbitrary when there is a mass and limited marrow involvement. This revision follows the convention used in many treatment protocols and suggests that when a mass is present and 25% or more of the nucleated cells in the bone marrow are lymphoblasts, a diagnosis of lymphoblastic leukemia is preferred over lymphoblastic lymphoma. Because ALL rarely presents with low BM blast counts, the diagnosis of ALL should be deferred if there are fewer than 20% blasts in the BM until there is definitive evidence to confirm the diagnosis. However, in the unusual case that a patient presents with less than 20% lymphoblasts in the BM and no evidence of an extramedullary mass, but demonstrates one of the known recurring cytogenetic abnormalities associated with ALL (see Table 2), the patient may be considered to have

lymphoblastic leukemia. However, the finding of less than 20% unequivocal lymphoblasts in the BM should also prompt a search for lymphoblastic lymphoma in an extramedullary location.

In the case of B lymphoblastic leukemia/lymphoma (a name change from the previous edition, in which “precursor B lymphoblastic leukemia/lymphoma” was used), numerous reports have demonstrated that recurring genetic abnormalities are associated with sufficiently unique clinical, immunophenotypic, and/or prognostic features so that they can be considered as distinct entities (Table 2).¹¹⁴⁻¹¹⁸ Although most of the specific chromosomal abnormalities may be detected by routine cytogenetic analysis, t(12;21)(p13;q22); *ETV6-RUNX1* is a submicroscopic rearrangement that is usually detected by FISH analysis. This abnormality is very important prognostically and it should always be searched for in childhood B-ALL, as should the *BCR-ABL1* in patients of any age, by routine cytogenetic analysis, FISH, or RT-PCR. Although rare, B lymphoblastic leukemia/lymphoma with t(5;14)(q31;q32); *IL3-IGH* should be considered in the differential diagnosis of peripheral blood eosinophilia. The eosinophils are reactive, and in some patients the percentage of blasts is relatively low in the bone marrow, in which case the chromosomal abnormality is sufficient to confirm the diagnosis. The finding of prominent eosinophilia in either B or T lymphoblastic leukemia/lymphoma should also prompt consideration of a rearrangement involving *FGFR1*, which, if present, would allow the case to be classified as lymphoblastic leukemia/lymphoma associated with an abnormality of *FGFR1*. If none of the specific genetic abnormalities listed in Table 2 are found, the designation of “B lymphoblastic leukemia/lymphoma, not otherwise specified,” is appropriate. It is important to understand that the term B-ALL should not be used for Burkitt lymphoma/leukemia, which is a neoplasm of mature B cells.

The neoplastic cells of 50% to 70% of patients with T lymphoblastic leukemia/lymphoma demonstrate abnormal karyotypes. The most common recurrent abnormalities are translocations that involve the alpha and delta T-cell receptor loci at 14q11.2, the beta locus at 7q35, or the gamma locus at 7p14-15 and that involve a growing list of partner genes.¹¹⁹ Although of pathogenetic significance, these abnormalities are not as clearly associated with unique biologic features as seen with some of the genetic abnormalities associated with B lymphoblastic leukemia/lymphoma, and thus the precursor T-cell neoplasms are not subdivided further according to their genetic defects.

Significant changes in the diagnosis and classification of precursor B- and T-cell neoplasms

1. The nomenclature has been changed from “precursor B lymphoblastic leukemia/lymphoma” and “precursor T lymphoblastic leukemia/lymphoma” to “B lymphoblastic leukemia/lymphoma” and “T lymphoblastic leukemia/lymphoma.”
2. B lymphoblastic leukemia/lymphoma is further divided into 7 distinct entities defined largely by specific recurring chromosomal abnormalities; cases of B-ALL lacking these abnormalities are considered as “not otherwise specified.”

Summary

The revised 2008 WHO classification of myeloid neoplasms and acute leukemia attempts to provide an up-to-date classification system that is based on recently published, peer-reviewed data. Already, in the few months since the publication of the 4th edition,

new information is accumulating that will eventually lead to our ability to recognize new diseases and change our criteria for the diseases already described. To continue to be useful in the clinics as well as in the evaluation of data from clinical trials and laboratory investigation, the classification must therefore be continually reviewed, updated, and tested. The exercise of developing the WHO document has produced a model of cooperation between pathologists, clinicians, and clinical scientists of all nationalities—a cooperation that must continue into the future for the benefit of patients with myeloid neoplasms and acute leukemia.

Acknowledgments

We thank Elizabeth Lanzl of the University of Chicago Editorial Office for assistance in copy editing the manuscript.

The Myeloid Clinical Advisory Committee (CAC) Meeting was supported through the generous contributions of the Leukemia Clinical Research Foundation, University of Chicago Cancer Research Center, Celgene Corporation, Genentech Incorporated, Pharmion Corporation, MGI Pharma, ARUP Laboratories, and Beckman-Coulter Corporation. No contributor was involved in the CAC program format, discussions, or conclusions.

Authorship

The WHO classification of myeloid neoplasms was developed and revised in part through discussions at the Myeloid Clinical Advisory Committee Meeting held in Chicago, Illinois, in February 2007. All authors of the manuscript served as session chairs and/or major presenters at the meeting and made major contributions to the classification described in this manuscript.

Contribution: J.W.V. drafted the initial manuscript; J.T. wrote and revised portions of the MPN section of the manuscript; D.A.A.

wrote and revised portions of the AML section of the manuscript; R.D.B. wrote and revised portions of the MDS section of the manuscript; M.J.B. wrote and revised portions of the ALL and MPAL sections of the manuscript; A.P. wrote and revised sections pertinent to phenotypic data; N.L.H. wrote and revised portions of the manuscript; M.M.L.B. wrote, reviewed, and revised genetic portions of manuscript; E.H.-L. contributed to the MDS classification and partially revised the manuscript; A.T. contributed to the MPN classification and partially revised the manuscript; and C.D.B. chaired the CAC meeting, contributed to all portions of classification, and helped write and revise the manuscript.

Conflict-of-interest disclosure: J.W.V. is a consultant with Celgene Corporation but has no conflicts of interest relevant to this manuscript. M.M.L.B. is a consultant for Baxter Health Care but has no conflicts of interest relevant to this manuscript. The remaining authors declare no competing financial interests.

Correspondence: James Vardiman, University of Chicago Hospitals, 5841 S Maryland Ave, MC0008, Chicago, IL 60637; e-mail: james.vardiman@uchospitals.edu.

Appendix

Members of the Myeloid and Acute Leukemia Clinical Advisory Committee (CAC) attending the CAC meeting, Chicago, IL, February 2007, included R. Arceci, Baltimore, MD; T. Barbui, Bergamo, Italy; J. Bennett, Rochester NY; C. D. Bloomfield (Meeting Chair), Columbus, OH; W. Carroll, New York, NY; T. M. de Witte, Nijmegen, The Netherlands; H. Dohner, Ulm, Germany; E. Estey, Seattle, WA; R. Foa, Rome, Italy; B. Falini, Perugia, Italy; P. Fenaux, Paris, France; D. G. Gilliland, Boston, MA; U. Germing, Dusseldorf, Germany; P. Greenberg, Stanford, CA; E. Hellström-Lindberg (Chair, MDS), Stockholm, Sweden; R. Larson, Chicago, IL; M. M. Le Beau, Chicago, IL; A. List, Tampa, FL; T. Naoe, Nagoya, Japan; C. M. Niemeyer, Freiburg, Germany; K. Shannon, San Francisco, CA; M. Tallman, Chicago, IL; A. Tefferi, Rochester MN (Chair, MPN)

References

- Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. World Health Organization Classification of Tumours. Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues. Lyon, France: IARC; 2001.
- Swerdlow SH, Campo E, Harris NL, et al, eds. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Lyon, France: IARC; 2008.
- Harris NL, Jaffe ES, Stein H, et al. A revised European-American classification of lymphoid neoplasms: a proposal from the International Lymphoma Study Group. *Blood*. 1994;84:1361-1392.
- Vardiman JW, Harris NL, Brunning RD. The World Health Organization (WHO) classification of the myeloid neoplasms. *Blood*. 2002;100:2292-2302.
- Mufti GJ, Bennett JM, Goasguen J, et al. Diagnosis and classification of myelodysplastic syndrome: International Working Group on Morphology of myelodysplastic syndrome (IWGM-MDS) consensus proposals for the definition and enumeration of myeloblasts and ring sideroblasts. *Haematologica*. 2008;93:1712-1717.
- Orazi A, Albitar M, Heerema NA, Haskins S, Neiman RS. Hypoplastic myelodysplastic syndromes can be distinguished from acquired aplastic anemia by CD34 and PCNA immunostaining of bone marrow biopsy specimens. *Am J Clin Pathol*. 1997;107:268-274.
- Cools J, DeAngelo DJ, Gotlib J, et al. A tyrosine kinase created by fusion of the PDGFRA and FIP1L1 genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *N Engl J Med*. 2003;348:1201-1214.
- James C, Ugo V, Le Couedic JP, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature*. 2005;434:1144-1148.
- Kralovics R, Passamonti F, Buser AS, et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med*. 2005;352:1779-1790.
- Levine RL, Wadleigh M, Cools J, et al. Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell*. 2005;7:387-397.
- Tefferi A, Lasho TL, Gilliland G. JAK2 mutations in myeloproliferative disorders [letter]. *N Engl J Med*. 2005;353:1416-1417, author reply 1416-1417.
- Pardanani AD, Levine RL, Lasho T, et al. MPL515 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood*. 2006;108:3472-3476.
- Pardanani A, Akin C, Valent P. Pathogenesis, clinical features, and treatment advances in mastocytosis. *Best Pract Res Clin Haematol*. 2006;19:595-615.
- Loh ML, Vattikuti S, Schubert S, et al. Mutations in PTPN11 implicate the SHP-2 phosphatase in leukemogenesis. *Blood*. 2004;103:2325-2331.
- Stephens K, Weaver M, Leppig KA, et al. Interstitial uniparental isodisomy at clustered breakpoint intervals is a frequent mechanism of NF1 inactivation in myeloid malignancies. *Blood*. 2006;108:1684-1689.
- Tartaglia M, Niemeyer CM, Fragale A, et al. Somatic mutations in PTPN11 in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia. *Nat Genet*. 2003;34:148-150.
- Willman CL. Molecular genetic features of myelodysplastic syndromes (MDS). *Leukemia*. 1998;12(suppl 1):S2-S6.
- Hirsch-Ginsberg C, LeMaistre AC, Kantarjian H, et al. RAS mutations are rare events in Philadelphia chromosome-negative/bcr gene rearrangement-negative chronic myelogenous leukemia, but are prevalent in chronic myelomonocytic leukemia. *Blood*. 1990;76:1214-1219.
- Padua RA, Carter G, Hughes D, et al. RAS mutations in myelodysplasia detected by amplification, oligonucleotide hybridization, and transformation. *Leukemia*. 1988;2:503-510.
- Mrozek K, Bloomfield CD. Chromosome aberrations, gene mutations and expression changes, and prognosis in adult acute myeloid leukemia. *Hematology Am Soc Hematol Educ Program*. 2006:169-177.
- Mrozek K, Marcucci G, Paschka P, Whitman SP, Bloomfield CD. Clinical relevance of mutations and gene-expression changes in adult acute myeloid leukemia with normal cytogenetics: are we

- ready for a prognostically prioritized molecular classification? *Blood*. 2007;109:431-448.
22. Paschka P, Marcucci G, Ruppert AS, et al. Adverse prognostic significance of KIT mutations in adult acute myeloid leukemia with inv(16) and t(8;21): a Cancer and Leukemia Group B Study. *J Clin Oncol*. 2006;24:3904-3911.
 23. Falini B, Mecucci C, Tiacci E, et al. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med*. 2005;352:254-266.
 24. Dohner K, Schlenk RF, Habdank M, et al. Mutant nucleophosmin (NPM1) predicts favorable prognosis in younger adults with acute myeloid leukemia and normal cytogenetics: interaction with other gene mutations. *Blood*. 2005;106:3740-3746.
 25. Greene ME, Mundschaug G, Wechsler J, et al. Mutations in GATA1 in both transient myeloproliferative disorder and acute megakaryoblastic leukemia of Down syndrome. *Blood Cells Mol Dis*. 2003;31:351-356.
 26. Levine RL, Gilliland DG. Myeloproliferative disorders. *Blood*. 2008;112:2190-2198.
 27. Jones AV, Kreil S, Zoi K, et al. Widespread occurrence of the JAK2 V617F mutation in chronic myeloproliferative disorders. *Blood*. 2005;106:2162-2168.
 28. Baxter EJ, Scott LM, Campbell PJ, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet*. 2005;365:1054-1061.
 29. Scott LM, Tong W, Levine RL, et al. JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med*. 2007;356:459-468.
 30. Thiele J, Kvasnicka HM, Orazi A. Bone marrow histopathology in myeloproliferative disorders—current diagnostic approach. *Semin Hematol*. 2005;42:184-195.
 31. Lengfelder E, Hochhaus A, Kronawitter U, et al. Should a platelet limit of $600 \times 10^9/l$ be used as a diagnostic criterion in essential thrombocythemia? An analysis of the natural course including early stages. *Br J Haematol*. 1998;100:15-23.
 32. Regev A, Stark P, Blickstein D, Lahav M. Thrombotic complications in essential thrombocythemia with relatively low platelet counts. *Am J Hematol*. 1997;56:168-172.
 33. Sacchi S, Vinci G, Gugliotta L, et al. Diagnosis of essential thrombocythemia at platelet counts between 400 and $600 \times 10^9/l$. Gruppo Italiano Malattie Mieloproliferative Croniche (GIMMC). *Haematologica*. 2000;85:492-495.
 34. Wilkins BS, Erber WN, Bareford D, et al. Bone marrow pathology in essential thrombocythemia: interobserver reliability and utility for identifying disease subtypes. *Blood*. 2008;111:60-70.
 35. Thiele J, Kvasnicka HM. Chronic myeloproliferative disorders with thrombocythemia: a comparative study of two classification systems (PVSG, WHO) on 839 patients. *Ann Hematol*. 2003;82:148-152.
 36. Florena AM, Tripodo C, Iannitto E, Porcasi R, Ingrao S, Franco V. Value of bone marrow biopsy in the diagnosis of essential thrombocythemia. *Haematologica*. 2004;89:911-919.
 37. Gianelli U, Vener C, Raviele PR, et al. Essential thrombocythemia or chronic idiopathic myelofibrosis? A single-center study based on hematopoietic bone marrow histology. *Leuk Lymphoma*. 2006;47:1774-1781.
 38. Golub TR, Barker GF, Lovett M, Gilliland DG. Fusion of PDGF receptor beta to a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell*. 1994;77:307-316.
 39. Keene P, Mendelow B, Pinto MR, et al. Abnormalities of chromosome 12p13 and malignant proliferation of eosinophils: a nonrandom association. *Br J Haematol*. 1987;67:25-31.
 40. Bain BJ, Fletcher SH. Chronic eosinophilic leukemias and the myeloproliferative variant of the hyper-eosinophilic syndrome. *Immunol Allergy Clin North Am*. 2007;27:377-388.
 41. Steer EJ, Cross NC. Myeloproliferative disorders with translocations of chromosome 5q31-35: role of the platelet-derived growth factor receptor Beta. *Acta Haematol*. 2002;107:113-122.
 42. Macdonald D, Reiter A, Cross NC. The 8p11 myeloproliferative syndrome: a distinct clinical entity caused by constitutive activation of FGFR1. *Acta Haematol*. 2002;107:101-107.
 43. Abruzzo LV, Jaffe ES, Cotelingam JD, Whang-Peng J, Del Duca V Jr, Medeiros LJ. T-cell lymphoblastic lymphoma with eosinophilia associated with subsequent myeloid malignancy. *Am J Surg Pathol*. 1992;16:236-245.
 44. Metzgeroth G, Walz C, Score J, et al. Recurrent finding of the FIP1L1-PDGFRFA fusion gene in eosinophilia-associated acute myeloid leukemia and lymphoblastic T-cell lymphoma. *Leukemia*. 2007;21:1183-1188.
 45. Nagata H, Worobec AS, Oh CK, et al. Identification of a point mutation in the catalytic domain of the protooncogene c-kit in peripheral blood mononuclear cells of patients who have mastocytosis with an associated hematologic disorder. *Proc Natl Acad Sci U S A*. 1995;92:10560-10564.
 46. Levine RL, Loriaux M, Huntly BJ, et al. The JAK2V617F activating mutation occurs in chronic myelomonocytic leukemia and acute myeloid leukemia, but not in acute lymphoblastic leukemia or chronic lymphocytic leukemia. *Blood*. 2005;106:3377-3379.
 47. Steensma DP, Dewald GW, Lasho TL, et al. The JAK2 V617F activating tyrosine kinase mutation is an infrequent event in both "atypical" myeloproliferative disorders and myelodysplastic syndromes. *Blood*. 2005;106:1207-1209.
 48. Sugimoto K, Hirano N, Toyoshima H, et al. Mutations of the p53 gene in myelodysplastic syndrome (MDS) and MDS-derived leukemia. *Blood*. 1993;81:3022-3026.
 49. Germing U, Gattermann N, Strupp C, Aivado M, Aul C. Validation of the WHO proposals for a new classification of primary myelodysplastic syndromes: a retrospective analysis of 1600 patients. *Leuk Res*. 2000;24:983-992.
 50. Germing U, Strupp C, Knipp S, et al. Chronic myelomonocytic leukemia in the light of the WHO proposals. *Haematologica*. 2007;92:974-977.
 51. Hellstrom-Lindberg E. Myelodysplastic syndromes: an historical perspective. *Hematology Am Soc Hematol Educ Program*. 2008:42.
 52. Raya JM, Arenillas L, Domingo A, et al. Refractory anemia with ringed sideroblasts associated with thrombocytosis: comparative analysis of marked with non-marked thrombocytosis, and relationship with JAK2 V617F mutational status. *Int J Hematol*. 2008;88:387-395.
 53. Boissinot M, Garand R, Hamidou M, Hermouet S. The JAK2-V617F mutation and essential thrombocythemia features in a subset of patients with refractory anemia with ring sideroblasts (RARS). *Blood*. 2006;108:1781-1782.
 54. Remacha AF, Nomdedeu JF, Puget G, et al. Occurrence of the JAK2 V617F mutation in the WHO provisional entity: myelodysplastic/myeloproliferative disease, unclassifiable-refractory anemia with ringed sideroblasts associated with marked thrombocytosis [letter]. *Haematologica*. 2006;91:719-720.
 55. Vardiman JW. Hematopathological concepts and controversies in the diagnosis and classification of myelodysplastic syndromes. *Hematology Am Soc Hematol Educ Program*. 2006;199-204.
 56. Bowen D, Culligan D, Jowitz S, et al. Guidelines for the diagnosis and therapy of adult myelodysplastic syndromes. *Br J Haematol*. 2003;120:187-200.
 57. Kennedy GA, Kay TD, Johnson DW, et al. Neutrophil dysplasia characterised by a pseudo-Pelger-Huet anomaly occurring with the use of mycophenolate mofetil and ganciclovir following renal transplantation: a report of five cases. *Pathology*. 2002;34:263-266.
 58. Brunning R, McKenna R, eds. Tumors of the Bone Marrow. Atlas of Tumor Pathology. Washington, DC: Armed Forces Institute of Pathology; 1994.
 59. Kouides PA, Bennett JM. Morphology and classification of the myelodysplastic syndromes and their pathologic variants. *Semin Hematol*. 1996;33:95-110.
 60. Bain BJ. The bone marrow aspirate of healthy subjects. *Br J Haematol*. 1996;94:206-209.
 61. Maciejewski JP, Rissitano A, Sloan EM, Nunez O, Young NS. Distinct clinical outcomes for cytogenetic abnormalities evolving from aplastic anemia. *Blood*. 2002;99:3129-3135.
 62. Steensma DP, Dewald GW, Hodnefield JM, Tefferi A, Hanson CA. Clonal cytogenetic abnormalities in bone marrow specimens without clear morphologic evidence of dysplasia: a form fruste of myelodysplasia? *Leuk Res*. 2003;27:235-242.
 63. Gupta V, Brooker C, Tooze JA, et al. Clinical relevance of cytogenetic abnormalities at diagnosis of acquired aplastic anaemia in adults. *Br J Haematol*. 2006;134:95-99.
 64. Pierre RV, Hoagland HC. Age-associated aneuploidy: loss of Y chromosome from human bone marrow cells with aging. *Cancer*. 1972;30:889-894.
 65. Della Porta MG, Malcovati L, Invernizzi R, et al. Flow cytometry evaluation of erythroid dysplasia in patients with myelodysplastic syndrome. *Leukemia*. 2006;20:549-555.
 66. Kussick SJ, Wood BL. Four-color flow cytometry identifies virtually all cytogenetically abnormal bone marrow samples in the workup of non-CML myeloproliferative disorders. *Am J Clin Pathol*. 2003;120:854-865.
 67. Stetler-Stevenson M, Arthur DC, Jabbour N, et al. Diagnostic utility of flow cytometric immunophenotyping in myelodysplastic syndrome. *Blood*. 2001;98:979-987.
 68. Wimazal F, Fonatsch C, Thalhammer R, et al. Idiopathic cytopenia of undetermined significance (ICUS) versus low risk MDS: the diagnostic interface. *Leuk Res*. 2007;31:1461-1468.
 69. Strupp C, Gattermann N, Giagounidis A, et al. Refractory anemia with excess of blasts in transformation: analysis of reclassification according to the WHO proposals. *Leuk Res*. 2003;27:397-404.
 70. Lambertenghi-Deliliers G, Orazi A, Luksch R, Annaloro C, Soligo D. Myelodysplastic syndrome with increased marrow fibrosis: a distinct clinicopathological entity. *Br J Haematol*. 1991;78:161-166.
 71. Tuzuner N, Cox C, Rowe JM, Watrous D, Bennett JM. Hypocellular myelodysplastic syndromes (MDS): new proposals. *Br J Haematol*. 1995;91:612-617.
 72. Steensma DP, Hanson CA, Letendre L, Tefferi A. Myelodysplasia with fibrosis: a distinct entity? *Leuk Res*. 2001;25:829-838.
 73. Elghetany MT, Hudnall SD, Gardner FH. Peripheral blood picture in primary hypocellular refractory anemia and idiopathic acquired aplastic anemia: an additional tool for differential diagnosis. *Haematologica*. 1997;82:21-24.
 74. Barrett J, Sauntharajah Y, Mouldrem J. Myelodysplastic syndrome and aplastic anemia: distinct entities or diseases linked by a common pathophysiology? *Semin Hematol*. 2000;37:15-29.
 75. Hasle H, Niemeyer CM, Chessells JM, et al. A pediatric approach to the WHO classification of myelodysplastic and myeloproliferative diseases. *Leukemia*. 2003;17:277-282.
 76. Kelly LM, Gilliland DG. Genetics of myeloid leukemias. *Annu Rev Genomics Hum Genet*. 2002;3:179-198.

77. Schlenk RF, Dohner K, Krauter J, et al. Mutations and treatment outcome in cytogenetically normal acute myeloid leukemia. *N Engl J Med*. 2008;358:1909-1918.
78. Thiede C, Koch S, Creutzig E, et al. Prevalence and prognostic impact of NPM1 mutations in 1485 adult patients with acute myeloid leukemia (AML). *Blood*. 2006;107:4011-4020.
79. Zelent A, Guidez F, Melnick A, Waxman S, Licht JD. Translocations of the RARalpha gene in acute promyelocytic leukemia. *Oncogene*. 2001;20:7186-7203.
80. Meyer C, Schneider B, Jakob S, et al. The MLL recombinome of acute leukemias. *Leukemia*. 2006;20:777-784.
81. Shih LY, Liang DC, Fu JF, et al. Characterization of fusion partner genes in 114 patients with de novo acute myeloid leukemia and MLL rearrangement. *Leukemia*. 2006;20:218-223.
82. Caligiuri MA, Strout MP, Lawrence D, et al. Rearrangement of ALL1 (MLL) in acute myeloid leukemia with normal cytogenetics. *Cancer Res*. 1998;58:55-59.
83. Bitter MA, Neilly ME, Le Beau MM, Pearson MG, Rowley JD. Rearrangements of chromosome 3 involving bands 3q21 and 3q26 are associated with normal or elevated platelet counts in acute nonlymphocytic leukemia. *Blood*. 1985;66:1362-1370.
84. Secker-Walker LM, Mehta A, Bain B. Abnormalities of 3q21 and 3q26 in myeloid malignancy: a United Kingdom Cancer Cytogenetic Group study. *Br J Haematol*. 1995;91:490-501.
85. Pearson MG, Vardiman JW, Le Beau MM, et al. Increased numbers of marrow basophils may be associated with a t(6;9) in ANLL. *Am J Hematol*. 1985;18:393-403.
86. Soupir CP, Vergilio JA, Dal Cin P, et al. Philadelphia chromosome-positive acute myeloid leukemia: a rare aggressive leukemia with clinicopathologic features distinct from chronic myeloid leukemia in myeloid blast crisis. *Am J Clin Pathol*. 2007;127:642-650.
87. Keung YK, Beaty M, Powell BL, Molnar I, Buss D, Pettenati M. Philadelphia chromosome positive myelodysplastic syndrome and acute myeloid leukemia-retrospective study and review of literature. *Leuk Res*. 2004;28:579-586.
88. Paietta E, Racevskis J, Bennett JM, et al. Biologic heterogeneity in Philadelphia chromosome-positive acute leukemia with myeloid morphology: the Eastern Cooperative Oncology Group experience. *Leukemia*. 1998;12:1881-1885.
89. Arber DA, Stein AS, Carter NH, Ikle D, Forman SJ, Slovak ML. Prognostic impact of acute myeloid leukemia classification. Importance of detection of recurring cytogenetic abnormalities and multilineage dysplasia on survival. *Am J Clin Pathol*. 2003;119:672-680.
90. Yanada M, Suzuki M, Kawashima K, et al. Long-term outcomes for unselected patients with acute myeloid leukemia categorized according to the World Health Organization classification: a single-center experience. *Eur J Haematol*. 2005;74:418-423.
91. Weinberg OK, Seetharam M, Ren L, et al. Clinical characterization of acute myeloid leukemia with myelodysplasia-related changes as defined by the 2008 WHO classification system. *Blood*. 2009;113:1906-1908.
92. Wandt H, Schakel U, Kroschinsky F, et al. MLD according to the WHO classification in AML has no correlation with age and no independent prognostic relevance as analyzed in 1766 patients. *Blood*. 2008;111:1855-1861.
93. Haferlach T, Schoch C, Loeffler H, et al. Morphologic dysplasia in de novo acute myeloid leukemia (AML) is related to unfavorable cytogenetics but has no independent prognostic relevance under the conditions of intensive induction therapy: results of a multiparameter analysis from the German AML Cooperative Group studies. *J Clin Oncol*. 2003;21:256-265.
94. Smith SM, Le Beau MM, Huo D, et al. Clinical-cytogenetic associations in 306 patients with therapy-related myelodysplasia and myeloid leukemia: the University of Chicago series. *Blood*. 2003;102:43-52.
95. Singh ZN, Huo D, Anastasi J, et al. Therapy-related myelodysplastic syndrome: morphologic subclassification may not be clinically relevant. *Am J Clin Pathol*. 2007;127:197-205.
96. Andersen MK, Larson RA, Mauritzson N, Schnittger S, Jhanwar SC, Pedersen-Bjerggaard J. Balanced chromosome abnormalities inv(16) and t(15;17) in therapy-related myelodysplastic syndromes and acute leukemia: report from an international workshop. *Genes Chromosomes Cancer*. 2002;33:395-400.
97. Bloomfield CD, Archer KJ, Mrozek K, et al. 11q23 balanced chromosome aberrations in treatment-related myelodysplastic syndromes and acute leukemia: report from an international workshop. *Genes Chromosomes Cancer*. 2002;33:362-378.
98. Rowley JD, Olney HJ. International workshop on the relationship of prior therapy to balanced chromosome aberrations in therapy-related myelodysplastic syndromes and acute leukemia: overview report. *Genes Chromosomes Cancer*. 2002;33:331-345.
99. Slovak ML, Bedell V, Popplewell L, Arber DA, Schoch C, Slater R. 21q22 balanced chromosome aberrations in therapy-related hematopoietic disorders: report from an international workshop. *Genes Chromosomes Cancer*. 2002;33:379-394.
100. Pileri SA, Ascani S, Cox MC, et al. Myeloid sarcoma: clinico-pathologic, phenotypic and cytogenetic analysis of 92 adult patients. *Leukemia*. 2007;21:340-350.
101. Gurbuxani S, Vyas P, Crispino JD. Recent insights into the mechanisms of myeloid leukemogenesis in Down syndrome. *Blood*. 2004;103:399-406.
102. Brink DS. Transient leukemia (transient myeloproliferative disorder, transient abnormal myelopoiesis) of Down syndrome. *Adv Anat Pathol*. 2006;13:256-262.
103. Massey GV, Zipursky A, Chang MN, et al. A prospective study of the natural history of transient leukemia (TL) in neonates with Down syndrome (DS): Children's Oncology Group (COG) study POG-9481. *Blood*. 2006;107:4606-4613.
104. Cella M, Jarrossay D, Facchetti F, et al. Plasmacytoid monocytes migrate to inflamed lymph nodes and produce large amounts of type I interferon. *Nat Med*. 1999;5:919-923.
105. Chaperot L, Bendriss N, Manches O, et al. Identification of a leukemic counterpart of the plasmacytoid dendritic cells. *Blood*. 2001;97:3210-3217.
106. Petrella T, Comeau MR, Maynadie M, et al. 'Agranular CD4+ CD56+ hematodermic neoplasm' (blastic NK-cell lymphoma) originates from a population of CD56+ precursor cells related to plasmacytoid monocytes. *Am J Surg Pathol*. 2002;26:852-862.
107. Piliichowska ME, Fleming MD, Pinkus JL, Pinkus GS. CD4+/CD56+ hematodermic neoplasm ("blastic natural killer cell lymphoma"): neoplastic cells express the immature dendritic cell marker BDCA-2 and produce interferon. *Am J Clin Pathol*. 2007;128:445-453.
108. Matutes E, Morilla R, Farahat N, et al. Definition of acute biphenotypic leukemia. *Haematologica*. 1997;82:64-66.
109. Hanson CA, Abaza M, Sheldon S, Ross CW, Schnitzer B, Stoolman LM. Acute biphenotypic leukaemia: immunophenotypic and cytogenetic analysis. *Br J Haematol*. 1993;84:49-60.
110. Sulak LE, Clare CN, Morale BA, Hansen KL, Montiel MM. Biphenotypic acute leukemia in adults. *Am J Clin Pathol*. 1990;94:54-58.
111. Campana D, Hansen-Hagge TE, Matutes E, et al. Phenotypic, genotypic, cytochemical, and ultrastructural characterization of acute undifferentiated leukemia. *Leukemia*. 1990;4:620-624.
112. Brito-Babapulle F, Pullon H, Layton DM, et al. Clinicopathological features of acute undifferentiated leukaemia with a stem cell phenotype. *Br J Haematol*. 1990;76:210-214.
113. Cuneo A, Ferrant A, Michaux JL, et al. Cytogenetic and clinicobiological features of acute leukemia with stem cell phenotype: study of nine cases. *Cancer Genet Cytogenet*. 1996;92:31-36.
114. Johansson B, Mertens F, Mitelman F. Clinical and biological importance of cytogenetic abnormalities in childhood and adult acute lymphoblastic leukemia. *Ann Med*. 2004;36:492-503.
115. Mitelman F. The Third International Workshop on Chromosomes in Leukemia. Lund, Sweden, July 21-25, 1980. Introduction. *Cancer Genet Cytogenet*. 1981;4:96-98.
116. Borkhardt A, Cazzaniga G, Viehmann S, et al. Incidence and clinical relevance of TEL/AML1 fusion genes in children with acute lymphoblastic leukemia enrolled in the German and Italian multicenter therapy trials. Associazione Italiana Ematologia Oncologia Pediatrica and the Berlin-Frankfurt-Munster Study Group. *Blood*. 1997;90:571-577.
117. Bloomfield CD, Goldman AI, Alimena G, et al. Chromosomal abnormalities identify high-risk and low-risk patients with acute lymphoblastic leukemia. *Blood*. 1986;67:415-420.
118. Pui CH, Robison LL, Look AT. Acute lymphoblastic leukaemia. *Lancet*. 2008;371:1030-1043.
119. Armstrong SA, Look AT. Molecular genetics of acute lymphoblastic leukemia. *J Clin Oncol*. 2005;23:6306-6315.



blood[®]

2009 114: 937-951
doi:10.1182/blood-2009-03-209262 originally published
online April 8, 2009

The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes

James W. Vardiman, Jüergen Thiele, Daniel A. Arber, Richard D. Brunning, Michael J. Borowitz, Anna Porwit, Nancy Lee Harris, Michelle M. Le Beau, Eva Hellström-Lindberg, Ayalew Tefferi and Clara D. Bloomfield

Updated information and services can be found at:
<http://www.bloodjournal.org/content/114/5/937.full.html>

Articles on similar topics can be found in the following Blood collections
[Free Research Articles](#) (5506 articles)
[Myeloid Neoplasia](#) (1996 articles)
[Perspectives](#) (223 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>