

Analysis of GPIIb/IIIa Receptor Number by Quantification of 7E3 Binding to Human Platelets

By Carrie L. Wagner, Mary A. Mascelli, Donald S. Neblock, Harlan F. Weisman, Barry S. Coller, and Robert E. Jordan

A large number of glycoprotein (GP) IIb/IIIa receptors are present on the surface of platelets. Studies to define precisely the number of GPIIb/IIIa receptors using specific monoclonal antibodies (MoAbs) or fibrinogen binding have, however, yielded varying estimates of receptor number. To refine the quantitative estimation of GPIIb/IIIa receptors on resting platelets, we have used the MoAb 7E3, which has high affinity for GPIIb/IIIa. Quantitative binding studies were performed using radiolabeled conjugates of 7E3 IgG, as well as fragments and derivatives of 7E3. For platelets obtained from any single individual, the numbers of 7E3 F(ab')₂ and IgG molecules bound per platelet were equivalent (~40,000), whereas the number of Fab molecules bound per platelet was consistently approximately twofold higher (~80,000). To investigate the basis of the quantitative disparity in bind-

ing of intact 7E3 and 7E3 F(ab')₂ versus 7E3 Fab, we studied the binding of a newly constructed, bispecific (Fab')₂ molecule containing only a single 7E3 combining site. Because this construct bound to the same extent as the Fab species, the larger size of the intact 7E3 and 7E3 F(ab')₂ molecules could not explain the reduced number of molecules that bound per platelet compared to the Fab fragment. Rather, it appears that the valency of the antibody is the critical factor determining the number of antibody molecules bound per platelet. Thus, we conclude that the binding of 7E3 Fab corresponds most closely with surface GPIIb/IIIa number and that the number of GPIIb/IIIa receptors is approximately 80,000 per platelet.

© 1996 by The American Society of Hematology.

THE PLATELET surface is densely populated with receptors for adhesive protein ligands.¹ Most prominent among these receptors is the glycoprotein IIb/IIIa complex (GPIIb/IIIa). An accurate assessment of the number of surface GPIIb/IIIa receptors is important for understanding the normal and pathological responsiveness of platelets. The GPIIb/IIIa receptor on resting platelets does not bind ligands with high affinity; after activation, the receptor develops a high binding affinity for fibrinogen and other adhesive proteins that enables the receptor to cross-link platelets one to another in stable platelet aggregates.² Approximately 40,000 fibrinogen molecules have been reported to bind per activated platelet,³⁻⁵ but this cannot be directly translated into receptor number because fibrinogen is dimeric and has a total of six regions that may potentially interact with GPIIb/IIIa receptors. Studies of the binding of small monovalent agents such as peptides and snake venom proteins⁶⁻⁸ to GPIIb/IIIa have yielded varying estimates of the number of molecules bound per platelet which may reflect differences in binding affinities or other properties. Monoclonal antibodies (MoAbs), by virtue of their high affinity and specificity for GPIIb/IIIa and the similarity of results using different antibodies, have been considered the most reliable indicators of GPIIb/IIIa copy number.

Regardless of the antibody used, a range of 35,000 to 50,000 antibody molecules have been reported to be bound per platelet,^{5,9-17} with approximately a twofold variability in the number of receptors present on platelets from different individuals.¹⁵ Our previous studies with antibody 7E3¹⁸ and studies by others using different anti-GPIIb/IIIa MoAbs^{19,20} indicated that more Fab' molecules bound per platelet than intact IgGs or F(ab')₂ fragments of the same antibodies. Various mechanisms can be proposed to explain these observations, among which are: (1) the existence of unique receptor pools that are accessible only to Fab fragments and not to IgGs, based on size-dependent access, (2) steric interference caused by the binding of another antibody molecule to an adjacent GPIIb/IIIa receptor, (3) internalization of Fab molecules into platelet α -granules, with perhaps binding to GPIIb/IIIa receptors in the α -granule membrane, (4) recruitment of GPIIb/IIIa receptors to the platelet surface from the α -granule pool triggered specifically by Fab fragments, or (5) the

simultaneous binding of the bivalent antibody species [IgG and F(ab')₂] to two different GPIIb/IIIa receptors. To differentiate between these mechanisms, we have studied the binding to platelets of 7E3 IgG, F(ab')₂, and Fab, as well as a newly constructed, bispecific (Fab')₂ molecule containing only a single 7E3 combining site, but having the same size as 7E3 F(ab')₂.

MATERIALS AND METHODS

Chemicals and reagents. All chemicals were purchased from commercial sources and used without purification.

Abbreviations. Purified antibody fragments of IgG are defined as follows: Fab denotes the IgG fragment produced by papain digestion, comprised of a light chain linked to the proteolytically cleaved heavy chain, and functionally monovalent for antigen binding; F(ab')₂ is a functionally bivalent fragment, a dimer of two Fab' fragments and is produced by pepsin digestion of IgG; Fab' is a monovalent fragment derived by mild reduction of the F(ab')₂ fragment followed by alkylation with iodoacetamide to prevent reoxidation of the sulfhydryl residues¹⁸; (Fab')₂ designates a bivalent fragment obtained by chemical or oxidative reconstitution of two Fab' fragments; a bispecific (Fab')₂ fragment is reconstituted from two different Fab' fragments.

MoAbs. The intact 7E3 and the 7E3 fragments bind to platelet GPIIb/IIIa and also to the α _v β ₃ vitronectin receptor. The murine 7E3,¹⁶ the murine/human chimeric 7E3,²¹ and OC125 (anti-ovarian cancer antigen)²² MoAbs were produced from hybridoma cell lines. The intact IgG molecules were isolated from cell supernatant by

From Centocor, Research and Development, Malvern, PA; and the Department of Medicine, Mount Sinai Medical Center, New York, NY.

Submitted December 18, 1995; accepted April 2, 1996.

Supported in part by Grant No. 19278 to B.S.C. from the National Heart, Lung and Blood Institute.

Address reprint requests to Carrie L. Wagner, PhD, 200 Great Valley Parkway, Malvern, PA 19355.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

© 1996 by The American Society of Hematology.

0006-4971/96/8803-0014\$3.00/0

affinity chromatography on Protein A-Sepharose (Pharmacia, Uppsala, Sweden). The Fab fragments were prepared by papain digestion and the F(ab')₂ fragments were prepared by pepsin digestion.²³ The fragments were further purified on Protein A-Sepharose to remove Fc-containing materials, and/or other gel filtration and/or ion exchange chromatography procedures to remove other contaminants. The OC125 Fab' fragment was prepared by mild reduction of the corresponding F(ab')₂ fragment followed by alkylation with iodoacetamide to prevent reoxidation of the disulfide bonds. The murine 7E3 Fab and the human/murine chimeric 7E3 Fab (ReoPro, abciximab; Centocor, Malvern, PA) fragments which are essentially identical in platelet binding and platelet function inhibition properties,²¹ were used interchangeably in these experiments, and are collectively identified as 7E3 Fab. The 7E3 IgG and the OC125 and 7E3 F(ab')₂ fragments used in these analyses were the native murine molecules.

Preparation of chemically cross-linked (Fab')₂ fragments. The method of preparation of the bispecific 7E3×OC125 (Fab')₂ fragment was the same used to prepare the bispecific 7E3×P4B6 (Fab')₂ fragment.²⁴ Briefly, parental F(ab')₂ fragments of an antibody to a cancer antigen not present on platelets, OC125, and 7E3 were reduced in 20 mmol/L L-cysteine and then the L-cysteine was removed by gel filtration. The reduced Fab' fragments were coupled using bis-maleimido methyl ether (BMME). Any remaining free sulfhydryl groups were capped with 5,5 dithiobis-(N, N')-nitrobenzoic acid, then the mixture was treated again with L-cysteine to reduce any recombined, non-BMME linked, parental (Fab')₂ fragments, and gel filtered using Sephacryl S-100 (Pharmacia) to remove the Fab' fragments. The bispecific species was resolved from the recombined, cross-linked parental molecules using hydrophobic interaction HPLC with a TSK phenyl-5-PW column (Bio-Rad, Hercules, CA). As a control, the BMME linking process was repeated using 7E3 Fab' for both arms of the recombined 7E3 (Fab')₂ molecule.

Estimation of antibody concentrations. Antibody concentrations (milligrams per milliliter) were assigned by dividing the absorption at 280 nm by 1.4.²⁵ Molecular weights were estimated for the 7E3 IgG, OC125 F(ab')₂, OC125 Fab', and the bispecific 7E3×OC125 (Fab')₂; and assigned based on the amino acid composition for the 7E3 Fab, 7E3 F(ab')₂ and the reconstituted 7E3 (Fab')₂ fragments. Molecular weights used were: 7E3 Fab = 47,600; 7E3 IgG = 150,000; OC125 F(ab')₂ = 100,000; OC125 Fab' = 50,000; bispecific 7E3×OC125 (Fab')₂ = 100,000; 7E3 F(ab')₂ = 97,600; reconstituted 7E3 (Fab')₂ = 97,600.

Radioiodination of proteins. Antibodies were iodinated using a modification of the Iodogen method²⁶ to a specific activity of 1 to 3 μCi/μg. Briefly, 90 μg of antibody was incubated with 0.5 mCi of Na¹²⁵I (Amersham Corp, Arlington Heights, IL) in tubes precoated with 5 μg of Iodogen (Pierce, Rockford, IL) for 5 minutes at room temperature. The reaction was halted by the addition of 5 mmol/L ascorbic acid and removal of the incubation mixture from the reaction vial. Free iodine was removed by gel filtration on a PD-10 column (Pharmacia). Antibodies were diluted with unlabeled antibody to achieve desired specific radioactivities.

Gel filtration chromatography. The monomeric nature of the iodinated antibody molecules and fragments was confirmed using gel filtration chromatography. The 7E3 Fab, F(ab')₂ and IgG molecules were subjected to a mock iodination using cold sodium iodide under comparable conditions used to radiolabel these same molecules. The gel filtration retention times of the cold I-labeled antibody and antibody fragments were analyzed using a Bio-Rad Bio-Sil SEC 250 column (300 × 7.8 mm) with a mobile phase of 0.2 mol/L sodium phosphate, pH. 6.8 at a flow rate of 1.0 mL per minute. The retention times of the unlabeled 7E3 Fab, 7E3 F(ab')₂ and 7E3 IgG molecules were identical to those of the corresponding I-labeled molecules (retention times of 9.64, 8.36, and 7.86 minutes, respec-

tively), indicating that none of these antibody molecules aggregated as a result of radioiodination.

Platelet preparation. Blood donors were selected from normal healthy adult volunteers who had not taken aspirin or an antihistaminic for a minimum of 10 days. Whole blood was drawn by venipuncture into 1:100 final volume of 40% trisodium citrate, and centrifuged at 600g for 6 minutes (or equivalent) at 22°C for preparation of platelet-rich plasma (PRP). For some experiments, platelets were isolated from the plasma by gel-filtration²⁷ of PRP on Sepharose 2B-Cl (Pharmacia) pre-equilibrated in modified Tyrode's buffer containing 20 mmol/L HEPES, pH 7.5, 187 mmol/L NaCl, 4 mmol/L KCl, 50 mmol/L Na₂HPO₄, 1 mmol/L MgCl₂, 5.5 mmol/L glucose, and 2% bovine serum albumin. PGE₁ (final concentration 1 nmol/L) was added to the gel-filtered platelet suspensions and the platelets were allowed to incubate at room temperature for at least 30 minutes before the initiation of binding studies. Platelet counts of the PRP and gel-filtered platelets were determined using a resistance particle counter (Coulter ZM, Hialeah, FL). Platelet counts of the PRP were adjusted to 300,000 ± 50,000 platelets/μL with autologous platelet-poor plasma if necessary.

Direct platelet binding assay. A direct binding method¹⁷ was used for measuring the number of antibody and antibody fragment molecules able to bind to platelets in both PRP and gel-filtered platelet suspensions. PRP was incubated with ¹²⁵I-labeled antibody or fragment for 30 to 60 minutes at room temperature. Replicate (3) 100-μL aliquots of the PRP were then centrifuged through 200 μL of 30% sucrose to separate free from platelet-bound antibody. The number of molecules of antibody bound per platelet was calculated as: the fraction of the total radioactivity that was platelet bound × the total antibody content (μg) × the appropriate conversion factor to obtain molecules per μg [Avogadro's number divided by (the antibody molecular weight × 10⁶ μg/g)], divided by the number of platelets in the sample.

Rate of dissociation of radiolabeled 7E3 Fab and F(ab')₂ from platelets. Studies were performed by first incubating a subsaturating amount of ¹²⁵I-7E3 Fab (0.2 μg/mL) or ¹²⁵I-7E3 F(ab')₂ (0.2 μg/mL) with PRP for 30 minutes at 37°C and determining the amount of radiolabeled antibody bound. Antibody dissociation was initiated by the addition of a 100-fold excess (20 μg/mL) of unlabeled 7E3 Fab or F(ab')₂ to the incubations containing ¹²⁵I-7E3 Fab or ¹²⁵I-7E3 F(ab')₂, respectively. The amount of bound radioactivity was determined following separation of the platelet-bound fraction from the unbound antibody fraction by centrifugation through 30% sucrose (described above). Platelet-bound radioactive antibody was determined at various time points up to 5 hours after the addition of unlabeled antibody and expressed as a percentage of the initial amount bound.

Sequential saturation platelet binding assay. A two-step assay was developed to assess the effect of prior exposure of platelets to various antibodies on the ability of ¹²⁵I-7E3 Fab to bind in a subsequent incubation step. PRP was first incubated with unlabeled antibodies at 10 μg/mL for 60 minutes at 22°C. In the second step, ¹²⁵I-7E3 Fab was added to a final concentration of 15 μg/mL, and incubated for an additional 30 minutes at 22°C. Quantification of bound radiolabeled antibody was performed as described above for the direct platelet binding assay.

Binding of antibody to thrombin-stimulated platelets. The binding of ¹²⁵I-7E3 IgG or ¹²⁵I-7E3 Fab to gel-filtered platelets was determined after stimulation with α-thrombin (Enzyme Research Laboratories, South Bend, IN), according to the procedure of Lombardo et al.²⁸ Aliquots of gel-filtered platelets were incubated with either 0.1 U/mL of thrombin or an equivalent volume of Tyrode's buffer for 10 minutes at 22 to 25°C. Hirudin (Sigma, St Louis, MO) was then added at a 16-fold excess (U/U) and allowed to incubate




Fig 1. Saturable binding of 7E3 IgG and 7E3 Fab to human platelets. Platelet-rich plasma (308,000 platelets/ μL) was prepared from the citrate-anticoagulated blood of a single donor, and incubated with varying concentrations of ^{125}I -7E3 IgG or ^{125}I -7E3 Fab for 60 minutes at 22°C . After separation of platelets from the plasma by centrifugation through 30% sucrose, the number of bound antibody molecules was calculated.

for 5 minutes at room temperature to neutralize the thrombin. Binding studies were then performed as described above.

Purification of platelet GPIIb/IIIa. GPIIb/IIIa was purified from outdated human platelets (50 to 100 U) obtained from the American Red Cross (Penn-Jersey Region, Philadelphia, PA). The platelets were pelleted, washed in 20 mmol/L Tris-Cl, 150 mmol/L NaCl, 0.4% sodium citrate, pH 7.0 buffer, and pelleted again. The GPIIb/IIIa was extracted by stirring overnight at room temperature in 1% Triton X-100, 20 mmol/L Tris-Cl, 150 mmol/L NaCl, 1 mmol/L CaCl_2 , 10^{-5} mol/L leupeptin, pH 7.4. The platelet fragments were separated from the soluble GPIIb/IIIa by centrifugation and filtration. The filtered supernatant was applied to an affinity column containing 10E5 (anti-GPIIb/IIIa MoAb) IgG. The bound GPIIb/IIIa was eluted with 50 mmol/L diethanolamine, 0.1% Triton X-100, 1 mmol/L CaCl_2 , pH 11.5. The eluted GPIIb/IIIa was immediately neutralized with 2 mol/L Tris-Cl, 10 mmol/L CaCl_2 , pH 7.0. The GPIIb/IIIa was concentrated in a stirred cell ultrafiltration device and stored with 0.1% sodium azide at 4°C . The GPIIb/IIIa purity and activity was assessed using nonreducing and reducing gradient sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), high-performance liquid chromatography gel filtration analysis, immunoblots, and binding of 7E3 and 10E5 antibodies in an enzyme-linked immunosorbent assay (ELISA). The isolated GPIIb/IIIa receptor exhibited approximately 98% purity by the gel filtration analysis,²⁹ with some minor contaminants observed by SDS-PAGE and immunoblot analyses.

GPIIb/IIIa binding ELISA. The binding of the native murine 7E3 $\text{F}(\text{ab}')_2$, the recombinant 7E3 $\text{F}(\text{ab}')_2$, and the bispecific 7E3 \times OC125 $\text{F}(\text{ab}')_2$ to purified GPIIb/IIIa was evaluated in a coated-plate ELISA.²⁴ The assay compared half-maximal binding concentrations of twofold dilutions of the 7E3 and OC125 fragments (evaluated from 1 to 1,000 ng/mL) to purified GPIIb/IIIa, which was detected using a peroxidase-conjugated, murine $\text{F}(\text{ab}')_2$ fragment-specific, goat antibody.

OC125 activity assay. The 7E3 $\text{F}(\text{ab}')_2$, OC125 $\text{F}(\text{ab}')_2$, OC125 $\text{F}(\text{ab}')_2$, and 7E3 \times OC125 $\text{F}(\text{ab}')_2$ molecules were analyzed for the ability to inhibit binding of OC125 IgG (anti-ovarian cancer antigen) in a CA 125 radioimmunoassay (RIA) (Centocor). The CA 125 RIA was a simultaneous, sandwich immunometric assay that used a single antibody, OC125, as both capture and signal antibody using anti-

body-coated beads and ^{125}I -labeled antibody.²² Over the concentration range evaluated (0.08 to 20 $\mu\text{mol/L}$), OC125 $\text{F}(\text{ab}')_2$ completely inhibited the binding of ^{125}I -labeled OC125. The 7E3 $\text{F}(\text{ab}')_2$ fragment had no inhibitory activity in the CA 125 RIA over the same concentration range. The bispecific 7E3 \times OC125 $\text{F}(\text{ab}')_2$ and the OC125 $\text{F}(\text{ab}')_2$ molecules attained complete inhibition of binding of the ^{125}I -labeled OC125 antibody at concentrations of 1.3 and 5 $\mu\text{mol/L}$, respectively.

RESULTS

Comparative binding of 7E3 IgG, 7E3 $\text{F}(\text{ab}')_2$, and 7E3 Fab. The results of an experiment in which the binding of ^{125}I -7E3 IgG and ^{125}I -7E3 Fab to unstimulated platelets in PRP measured after 1 hour of incubation at 22°C are shown in Fig 1. One hour exceeds the amount of time required to achieve equilibrium binding of all species of the 7E3 antibody. These results are representative of a number of similar experiments, and indicate that saturation binding was achieved for both forms of the 7E3 antibody after 1 hour at concentrations greater than or equal to 5 $\mu\text{g/mL}$. Approximately twice as many 7E3 Fab fragments were bound per platelet compared to 7E3 IgG ($\sim 80,000$ v $\sim 40,000$, respectively), which is consistent with our previous observations.¹⁸ Attempts to further refine the estimates of maximum binding by curve fitting, Scatchard analysis, or other linearizations of binding data did not significantly alter the estimates of maximum binding obtained by direct examination of the full binding curve data. Because saturation binding of both forms of 7E3 required a minimum concentration of 5 $\mu\text{g/mL}$, in subsequent studies antibody concentrations of ≥ 15 $\mu\text{g/mL}$ were routinely used.

Studies comparing the binding of 7E3 IgG and 7E3 Fab at a saturating concentration were performed in experiments with platelets obtained from 12 normal human donors (Fig 2). The inter-subject range of 7E3 IgG binding was 35,200 to 50,400 molecules per platelet and the range of 7E3 Fab




Fig 2. Individual quantitative measurements of ^{125}I -7E3 IgG and ^{125}I -7E3 Fab binding to platelets from 12 normal human donors. Platelets in PRP were incubated with saturating concentrations of either ^{125}I -7E3 Fab (15 $\mu\text{g/mL}$) or ^{125}I -7E3 IgG (18 $\mu\text{g/mL}$) for 30 minutes at 22°C . The lines connect the Fab and IgG binding values for each individual to highlight the consistent twofold difference in the extent of binding. For this population, the mean \pm SD number of 7E3 binding sites per platelet was $87,200 \pm 9,600$ for 7E3 Fab and $41,300 \pm 4,100$ for 7E3 IgG.




Fig 3. Binding measurements of ^{125}I -7E3 IgG, ^{125}I -7E3 F(ab')₂ and ^{125}I -7E3 Fab to platelets from seven normal donors. PRP was prepared from citrate-anticoagulated blood, and incubated with 15 $\mu\text{g}/\text{mL}$ of radiolabeled antibody for 60 minutes at 22°C. After separation of platelets from the plasma by centrifugation through 30% sucrose, the number of bound antibody molecules was calculated. The height of the bars indicates the average number of bound antibody molecules per platelet and the error bars correspond to 1 SD.

binding was 71,700 to 102,000 molecules per platelet. In this analysis, an average of 41,300 ($\pm 4,100$) molecules of 7E3 IgG bound per platelet, and 87,200 ($\pm 9,800$) 7E3 Fab molecules bound per platelet. Despite the variability in binding between individuals, the results for each patient showed a consistent near 2 to 1 relationship between Fab and IgG binding (mean \pm SD, 2.1 ± 0.1 ; range, 1.9 to 2.3).

The comparative binding of 7E3 IgG, 7E3 F(ab')₂, and 7E3 Fab to platelets from a panel of normal donors ($n = 7$) is presented in Fig 3. The mean \pm SD binding of 7E3 Fab was $81,900 \pm 9,200$ molecules per platelet. The numbers of 7E3 IgG and 7E3 F(ab')₂ molecules bound per platelet were similar ($40,600 \pm 4,100$ and $43,400 \pm 5,900$, respectively), and both were approximately 50% of the number of 7E3 Fab molecules bound per platelet. The similarity of 7E3 IgG and 7E3 F(ab')₂ binding suggests that the loss of the Fc domain of intact 7E3 IgG does not affect maximal binding, even though it reduces the molecular weight by approximately one third.

To further investigate the difference in binding of 7E3 F(ab')₂ and 7E3 Fab to platelets, the rates of dissociation of each fragment were compared. Dissociation of platelet-bound, ^{125}I -7E3 F(ab')₂ and ^{125}I -7E3 Fab was measured after addition of a 100-fold excess of the respective unlabeled antibody as shown in Fig 4. The ^{125}I -7E3 F(ab')₂ dissociated from platelets much slower than did ^{125}I -7E3 Fab under these conditions. After 5 hours of incubation with the competing unlabeled 7E3 F(ab')₂, 96% of the ^{125}I -7E3 F(ab')₂ remained bound to platelets. Only 42% of the ^{125}I -7E3 Fab remained bound after the same interval. The significantly slower dissociation of 7E3 F(ab')₂ is consistent with it binding to two different GPIIb/IIIa receptors simultaneously.

Studies to assess whether 7E3 Fab can bind to a subpopulation of GPIIb/IIIa receptors inaccessible to 7E3 IgG and 7E3 F(ab')₂. To assess whether all forms of 7E3 have equal accessibility to the same population(s) of GPIIb/IIIa recep-

tors, a two-step binding experiment was designed. If the larger, bivalent 7E3 molecules could not access certain GPIIb/IIIa receptors that the smaller 7E3 Fab fragments could engage, then radiolabeled 7E3 Fab should be able to bind to platelets pretreated with 7E3 IgG. In the first step, aliquots of PRP were preincubated for 1 hour with 10 $\mu\text{g}/\text{mL}$ of 7E3 Fab, 7E3 F(ab')₂, 7E3 IgG, or a non-platelet-specific control antibody, OC125 F(ab')₂. In the second step, ^{125}I -7E3 Fab (15 $\mu\text{g}/\text{mL}$) was added, and the incubation was continued for an additional 30 minutes. The rates of dissociation of 7E3 IgG¹⁸ and 7E3 F(ab')₂ (Fig 4) from the platelet surface are slow, ensuring that antibody that bound to the platelets in the preincubation step would not dissociate significantly in the subsequent 30-minute incubation step.

The results shown in Fig 5 indicate that preincubating platelets with 7E3 IgG or 7E3 F(ab')₂ reduced the subsequent binding of 7E3 Fab by 88% and 95%, respectively. Preincubation with unlabeled 7E3 Fab itself produced a similar reduction (94%) in radiolabeled 7E3 Fab binding. The control antibody, OC125 F(ab')₂, did not reduce 7E3 Fab binding. If the twofold increase in 7E3 Fab binding compared with 7E3 IgG or 7E3 F(ab')₂ binding were caused by differences in access to a subpopulation of GPIIb/IIIa receptors, one would have expected the 7E3 IgG and 7E3 F(ab')₂ preincubation to only decrease 7E3 Fab binding by approximately 50%.

Internal pools of GPIIb/IIIa receptors. Platelets contain an internal reservoir of GPIIb/IIIa receptors³⁰ that can join the outer platelet membrane surface when the platelet undergoes the release reaction and α -granule membranes fuse with the open canalicular system and ultimately the plasma mem-




Fig 4. Dissociation of 7E3 Fab and 7E3 F(ab')₂ from platelets. A subsaturating amount of ^{125}I -7E3 Fab (0.2 $\mu\text{g}/\text{mL}$) or ^{125}I -7E3 F(ab')₂ (0.2 $\mu\text{g}/\text{mL}$) was added to PRP and incubated for 30 minutes at 37°C in a 5% CO₂ incubator. The dissociation of ^{125}I -7E3 Fab or ^{125}I -7E3 F(ab')₂ was evaluated after addition of a 100-fold excess (20 $\mu\text{g}/\text{mL}$) of the respective unlabeled antibody fragment to the platelet suspension and removal of aliquots at the indicated time. The amount of residual bound radioactivity was determined after separation of the platelet-bound fraction from the unbound antibody by centrifugation through 30% sucrose, and expressed as a percentage of the initial amount bound.




Fig 5. Effect of prior exposure of platelets to saturating concentrations (10 $\mu\text{g}/\text{mL}$), of unlabeled 7E3 IgG, 7E3 F(ab')₂, 7E3 Fab or the control antibody OC125 F(ab')₂ on the subsequent binding of ¹²⁵I-7E3 Fab (15 $\mu\text{g}/\text{mL}$). Initial incubations were performed for 60 minutes at 22°C and the secondary incubation for 30 minutes at 22°C. After separation of platelets from the plasma, the number of platelet bound ¹²⁵I-7E3 molecules were quantified. The number of ¹²⁵I-7E3 Fab molecules is expressed as a percent of the binding that was measured in the absence of antibody in the initial incubation. The height of bars represents the average of the determinations from four normal donors and the error bars correspond to 1 SD.

brane.³¹ We considered the possibility that the 7E3 Fab fragment could uniquely bind to internal GPIIb/IIIa receptors under the conditions used in these binding studies, thus leading to a larger number of bound 7E3 Fab molecules per platelet. This could occur if 7E3 Fab caused externalization of these receptors or if the 7E3 Fab fragment was itself rapidly internalized and bound to internal receptors. To test the first possibility, we measured the release of the contents of both dense bodies (serotonin, adenosine triphosphate) and α -granules (β -thromboglobulin) when platelets were treated with 7E3 Fab, and it did not produce granule release (data not shown).

If 7E3 Fab, but not 7E3 IgG, was able to gain access to the internal store of receptors on unactivated platelets under the conditions of these binding experiments, then aliquots of activated and unactivated platelets incubated with ¹²⁵I-7E3 Fab should show little or no difference in the number of molecules of 7E3 Fab bound. In contrast, if the lower number of 7E3 IgG molecules bound per platelet were

caused by an inability to access internal GPIIb/IIIa receptor stores, then the activated platelet sample would be expected to show an increased binding of the ¹²⁵I-7E3 IgG relative to the unactivated platelet aliquot. To avoid platelet aggregation and fibrin formation by thrombin, studies were performed with gel-filtered platelets. Activation of the gel-filtered platelets with α -thrombin before incubation with radiolabeled 7E3 Fab or IgG significantly increased the number of both 7E3 IgG and 7E3 Fab molecules bound per platelet (1.4-fold and 1.3-fold, respectively) relative to the unactivated samples (Table 1). Thus, these data do not support either of the hypotheses that the twofold increase in 7E3 Fab binding compared to 7E3 IgG is due to unique activation of platelets by 7E3 Fab or to unique access of 7E3 Fab to the internal pool of GPIIb/IIIa present in unactivated platelets.

Studies of the effect of molecular size and valency using the bispecific 7E3 \times OC125 (Fab')₂ molecule. A bispecific antibody with the size of an F(ab')₂ fragment but with only a single binding site for GPIIb/IIIa was synthesized. To ensure that the activity of the chemically cross-linked bispecific (Fab')₂ molecule was comparable to the native 7E3 F(ab')₂, it was evaluated in a GPIIb/IIIa binding ELISA. The native 7E3 F(ab')₂, the recombinant 7E3 (Fab')₂, and the bispecific 7E3 \times OC125 (Fab')₂ had comparable half-maximal GPIIb/IIIa binding concentrations, whereas the native OC125 F(ab')₂ had no activity over the concentration range tested (data not shown). If antibody size is the major factor defining the number of 7E3 molecules bound, then the binding of the 7E3 \times OC125 (Fab')₂ should parallel that of the 7E3 F(ab')₂. If valency is the determining factor, then the bispecific (Fab')₂ molecule should bind to the same extent as 7E3 Fab, since the OC125 combining site does not recognize GPIIb/IIIa.

The binding of the bispecific 7E3 \times OC125 (Fab')₂ antibody and a control, reconstituted bivalent 7E3 (Fab')₂ fragment were tested against the platelets obtained from the same 7 donors used for the experiment described in Fig 3. As shown in Fig 6, the binding of the bispecific (Fab')₂ (76,800 \pm 6,300 molecules per platelet) was similar to the binding of the 7E3 Fab (81,000 \pm 9,200 molecules per platelet) (Fig 3), and both were significantly greater than the binding of native 7E3 F(ab')₂ (43,400 \pm 5,900 molecules per platelet) (Fig 3) or the reconstituted 7E3 (Fab')₂ (38,500 \pm 4,200) (Fig 6). The similarity in binding properties of the native 7E3 F(ab')₂ and the reconstituted 7E3 (Fab')₂ indicates that

Table 1. 7E3 Binding to Unstimulated and Thrombin-Stimulated Platelets

	Molecules Bound per Platelet			Ratio of No. of Molecules Bound for Stimulated Platelets/Unstimulated Platelets
	PRP	Unstimulated GF Platelets	Stimulated GF Platelets + Thrombin	
7E3 Fab	84,300 \pm 10,700	92,900 \pm 13,100	126,000 \pm 20,600	1.4 \pm 0.1
7E3 IgG	37,400 \pm 4,500	42,700 \pm 3,300	54,300 \pm 3,400	1.3 \pm 0.1

The number of radiolabeled 7E3 IgG or 7E3 Fab molecules bound per platelet was quantified using platelets from normal donors in both PRP and after gel filtration to remove fibrinogen. The binding of ¹²⁵I-7E3 IgG or ¹²⁵I-7E3 Fab was also measured using gel-filtered (GF) platelets after the platelets were stimulated for 10 minutes with thrombin (0.1 U/mL). All platelet samples were then incubated with the radioiodinated forms of 7E3 for 30 minutes at room temperature. After the separation of platelets from the unbound 7E3 molecules by centrifugation through 30% sucrose, the number of platelet-bound ¹²⁵I-7E3 molecules were quantified. The numbers represent the average of 5 determinations \pm 1 SD.




Fig 6. Binding of radiolabeled conjugates of the recombinant 7E3 (Fab')₂, the bispecific 7E3 × OC125 (Fab')₂ and OC125 F(ab')₂ to platelets. Platelets were obtained from the same seven normal donors shown in Fig 3. PRP was prepared from citrate-anticoagulated blood, and incubated with 15 μg/mL of radiolabeled antibody for 60 minutes at 22°C. After separation of platelets from the plasma by centrifugation through 30% sucrose, the number of bound antibody molecules was calculated. The height of the bars indicate the average number of molecules of each antibody species bound per platelet and the error bars correspond to 1 SD.

the BMME linkage procedure does not significantly alter the binding properties of 7E3. The nonspecific binding of radiolabeled OC125 F(ab')₂ control antibody fragment was 400 ± 200 molecules per platelet. Thus, the results with the bispecific 7E3×OC125 (Fab')₂ fragment indicate that the valency rather than the size of the 7E3 species determines the extent of binding to platelets.

DISCUSSION

Investigations designed to establish the number of cellular antigens or receptors routinely rely on quantitative determinations of ligand or specific MoAb binding. Quantitative evaluation of the binding of the ligand fibrinogen to the platelet as a method of estimating the number of GPIIb/IIIa receptors on the surface of unactivated platelets has serious limitations because (1) activation stimuli may not result in all GPIIb/IIIa receptors undergoing the transition to the high affinity ligand binding state, (2) the binding of one fibrinogen molecule may result in steric hindrance that limits the binding of other fibrinogen molecules, (3) internal pools of GPIIb/IIIa may join the surface, and (4) the stoichiometry of fibrinogen binding to GPIIb/IIIa may not be 1:1.

MoAbs and fragments of MoAbs that bind to the unactivated form of GPIIb/IIIa should, in theory, provide a more direct means for assessing receptor number on unactivated platelets, but issues of steric hindrance, access to internal stores, and the stoichiometry of binding may also affect the results. The consistent observation by us¹⁸ and others^{19,20} of a larger number of Fab or Fab' fragments binding per platelet than intact IgG molecules requires a rigorous examination if one is to obtain an accurate estimate of the number of platelet surface GPIIb/IIIa, and that is why the current studies were undertaken. We exploited the high-affinity binding

properties of the 7E3 MoAb to GPIIb/IIIa receptors on both unactivated and activated platelets, and the ability to make a variety of forms of the 7E3 antibody differing in size and valency. Although the rate of 7E3 F(ab')₂ and 7E3 Fab fragment binding to unactivated platelets is more rapid than observed for the larger IgG molecule,¹⁸ maximal platelet binding for all 7E3 molecules is obtained rapidly at concentrations of 5 μg/mL. For the present studies, binding incubations were performed for 30 to 60 minutes to ensure that equilibrium binding was attained. Previous studies established that the equilibrium dissociation constants for the interaction of 7E3 IgG, F(ab')₂ and Fab to resting platelets are similar at 4 nmol/L, 2 nmol/L, and 5 nmol/L, respectively (data not shown). Except for the experiments that required gel-filtered platelets, all other binding studies were conducted in PRP that was prepared from freshly obtained, citrate-anticoagulated blood from donors who had not taken antiplatelet agents.

In addition to platelet GPIIb/IIIa, 7E3 also binds to another integrin receptor, α_vβ₃, that also contains the GPIIIa (β₃) subunit. Although α_vβ₃ is present on platelets, the number of α_vβ₃ receptors is very small (~50 to 200 per platelet) representing less than 0.5% of the number of GPIIb/IIIa receptors.³² Thus, the potential contribution of platelet α_vβ₃ to total 7E3 binding was disregarded for these quantitative binding studies.

The results of this study indicate that the most plausible explanation for the twofold higher binding of 7E3 Fab compared with 7E3 IgG is that each of the two antibody-combining sites of the 7E3 IgG engages a separate and adjacent GPIIb/IIIa molecule. Several lines of evidence are consistent with this explanation. No evidence was found to support the existence of discrete surface GPIIb/IIIa receptor populations that are accessible only to the Fab fragment and not to F(ab')₂ or IgG. Thus, exposure of platelets to saturating concentrations of unlabeled 7E3 IgG or F(ab')₂ blocked the access of 7E3 Fab to the platelet surface nearly as well as did unlabeled 7E3 Fab. No evidence was obtained to suggest that the 7E3 Fab fragment produced release of α-granule contents or increased surface GPIIb/IIIa expression. Moreover, it is unlikely that selective internalization of 7E3 Fab accounts for the differences in binding because the number of molecules of 7E3 Fab that bound to platelets increased when platelets were preactivated with thrombin, to approximately the same extent as did platelets reacted with 7E3 IgG. It is also unlikely that the larger molecular size of the 7E3 IgG or F(ab')₂ fragment is responsible for limiting the extent of binding to one half the level of the 7E3 Fab fragment because the bispecific 7E3×OC125 (Fab')₂ fragment, containing only one 7E3 combining site, bound to platelets to the same extent as 7E3 Fab, ie, twice the level of 7E3 (Fab')₂. This result establishes that size alone is not a major determinant of the extent of binding and points, instead, to a bivalent mode of binding of IgG and F(ab')₂ fragments to surface GPIIb/IIIa as explaining the twofold disparity in binding. The results are most consistent with a bivalent binding of 7E3 IgG or F(ab')₂ to two adjacent receptors on the same platelet, because IgGs are ineffective at bridging between platelets³³ and no evidence of agglutination was

ever observed after the addition of 7E3 IgG or 7E3 F(ab')₂ to platelets. The extraordinary density of GPIIb/IIIa on the surface of platelets, with molecules estimated to be spaced approximately 160 Å apart,* adds plausibility to the hypothesis that the bivalent antibody molecules can actually engage two different receptors, even without having to postulate extensive movement of receptors in the membrane, because the two combining sites of a murine IgG₁ MoAbs have been estimated to be 117 to 134 Å apart.³⁴

Several laboratories have used MoAb-based techniques to provide estimates of approximately 40,000 GPIIb/IIIa receptors per platelet. This number is widely accepted, in part, because of its concordance with the number of fibrinogen molecules reported to bind to activated platelets.³⁴ Studies with another of our antibodies, 7H2, which is directed at GPIIIa, also showed a twofold increase in binding of 7H2 Fab compared to 7H2 IgG, with approximately 80,000 Fab molecules binding per platelet.²⁰ If the actual number of GPIIb/IIIa receptors is ~80,000, or twice the number of fibrinogen molecules that bind at saturation, this raises the interesting possibility that a single fibrinogen molecule may also bind to two different GPIIb/IIIa receptors on a single platelet. Each intact fibrinogen molecule contains two γ -chain dodecapeptide sequences that have been implicated in binding to GPIIb/IIIa and promoting platelet aggregation.³⁵ Additionally, fibrinogen contains four arginine-glycine-aspartic acid (RGD) sequences that may contribute alternative sites for GPIIb/IIIa binding, although current data would suggest that these RGD sites are less important than the dodecapeptide sequences.³⁶ Also uncertain is the degree to which fibrinogen is structurally constrained to minimize "cis" interactions with GPIIb/IIIa receptors on the surface of the same platelet as opposed to "trans" interactions with GPIIb/IIIa receptors on different platelets.³⁷ Nevertheless, the multivalent nature of the fibrinogen molecule suggests that at least bivalent binding to a single platelet is possible and may at least partly explain why ~40,000 fibrinogen molecules bind per platelet, even if there are twice as many GPIIb/IIIa receptors.

ACKNOWLEDGMENT

We thank Ellen Lance and Michael Spitz for their excellent technical assistance. We are also very grateful to Greg Brink for his graphics support and Joani Wendel for her typographical assistance.

REFERENCES

1. Phillips DR, Charo IF, Parise LV, Fitzgerald LA: The platelet membrane glycoprotein IIb-IIIa complex. *Blood* 71:831, 1988
2. Collier BS: Platelets in cardiovascular thrombosis and thrombolysis, in Fozzard HA (ed): *The Heart and Cardiovascular System* (ed 2). New York, NY, Raven, 1992, p 219

*The surface area of a platelet is estimated to be approximately 20 square microns (equivalent to $2 \times 10^9 \text{ \AA}^2$). If the surface area of the platelet is represented as a grid containing 80,000 squares each containing one GPIIb/IIIa receptor, then the area of each square would equal $2.5 \times 10^4 \text{ \AA}^2$. The distance between the midpoints of two adjacent squares (equivalent to the length of the side of each square) would then be estimated as 158 \AA ($\sqrt{2.5 \times 10^4 \text{ \AA}^2}$).

3. Bennett JS, Vilaire G: Exposure of platelet fibrinogen receptors by ADP or epinephrine. *J Clin Invest* 64:393, 1979
4. Marguerie GA, Edgington TS, Plow EF: Interaction of fibrinogen with its platelet receptor as part of a multistep reaction in ADP-mediated platelet aggregation. *J Biol Chem* 255:154, 1980
5. Niiya K, Hodson E, Bader R, Byers-Ward V, Koziol JA, Plow EF, Ruggeri ZM: Increased surface expression of the membrane GPIIb/IIIa complex induced by platelet activation. Relationship to the binding of fibrinogen and platelet aggregation. *Blood* 70:475, 1987
6. Wencel-Drake JD, Frelinger AL, Dieter MG, Lam SC-T: Arg-Gly-Asp-dependent occupancy of GPIIb/IIIa by Applagin: Evidence for internalization and cycling of a platelet integrin. *Blood* 81:62, 1993
7. Savage B, Marzec U, Chao B, Harker L, Maraganore J, Ruggeri Z: Binding of the snake venom-derived proteins Applagin and Echistatin to the arginine-glycine-aspartic acid recognition site(s) on platelet glycoprotein IIb/IIIa complex inhibits receptor function. *J Biol Chem* 265:11766, 1990
8. Tsao P, Bozarth J, Jackson S, Forsythe M, Flint S, Mousa S: Platelet GPIIb/IIIa receptor occupancy studies using a novel fluorescent cyclic Arg-Gly-Asp peptide. *Thrombosis Res* 77:543, 1995
9. Newman PJ, Allen RW, Kahn RA, Kunicki TJ: Quantitation of membrane GPIIIa on intact human platelets using the monoclonal antibody, AP-3. *Blood* 65:227, 1985
10. Tomiyama Y, Tsubakio T, Piotrowicz RS, Kurata Y, Loftus JC, Kunicki TJ: The Arg-Gly-Asp (RGD) recognition site of platelet glycoprotein IIb-IIIa on nonactivated platelets is accessible to high-affinity macromolecules. *Blood* 79:2303, 1992
11. Pidard D, Montgomery RR, Bennett JS, Kunicki TJ: Interaction of AP-2, a monoclonal antibody specific for the human platelet glycoprotein IIb-IIIa complex, with intact platelets. *J Biol Chem* 258:12582, 1983
12. McEver RP, Bennett EM, Martin MN: Identification of two structurally and functionally distinct sites on human platelet membrane glycoprotein IIb-IIIa using monoclonal antibodies. *J Biol Chem* 258:5269, 1983
13. Bennett JS, Hoxie JA, Leitman SF, Vilaire G, Cines DB: Inhibition of fibrinogen binding to stimulated human platelets by a monoclonal antibody. *Proc Natl Acad Sci USA* 80:2417, 1983
14. Ramsamooj P, Doellgast GJ, Hantgan RR: Inhibition of fibrin(ogen) binding to stimulated platelets by a monoclonal antibody specific for a conformational determinant of GPIIIa. *Thromb Res* 58:577, 1990
15. McEver RP, Baenziger NL, Majerus PW: Isolation and quantitation of the platelet membrane glycoprotein deficient in thrombasthenia using a monoclonal hybridoma antibody. *J Clin Invest* 66:1311, 1980
16. Collier BS, Peerschke EI, Scudder LE, Sullivan CA: A murine monoclonal antibody that completely blocks the binding of fibrinogen to platelets produces a thrombasthenic-like state in normal platelets and binds to glycoproteins IIb and/or IIIa. *J Clin Invest* 73:325, 1983
17. Collier BS: A new murine monoclonal antibody reports an activation-dependent change in the conformation and/or microenvironment of the platelet glycoprotein IIb/IIIa complex. *J Clin Invest* 76:101, 1985
18. Collier BS: Activation affects access to the platelet receptor for adhesive glycoproteins. *J Cell Biol* 103:451, 1986
19. Woods VL, Wolff LE, Keller DM: Resting platelets contain a substantial centrally located pool of GPIIb/IIIa complex which may be accessible to some but not other extracellular proteins. *J Biol Chem* 261:15242, 1986
20. Kutok JL, Collier BS: Partial inhibition of platelet aggregation

and fibrinogen binding by a murine monoclonal antibody to GPIIIa: Requirement for antibody bivalency. *Thromb Haemost* 72:964, 1994

21. Knight DM, Wagner C, Jordan R, McAleer MF, Derita R, Fass DN, Collier BS, Weisman H, Ghayeb J: Humanizing the constant region of the 7E3 murine monoclonal Fab antibody fragment reduces the immunogenicity of the variable region in humans. *Mol Immunol* 32:1271, 1995

22. Bast RC, Feeney M, Lazarus H, Nadler LM, Colvin RB, Knapp RC: Reactivity of a monoclonal antibody with human ovarian carcinoma. *J Clin Invest* 68:1331, 1981

23. Parham P: Preparation and purification of active fragments from mouse monoclonal antibodies, in Weir DM (ed): *Handbook of Experimental Immunocytochemistry: Immunocytochemistry*. London, UK, Blackwell Scientific, 1986 p 14.1

24. Neblock DS, Chang C-H, Mascelli MA, Fleek M, Stumpo L, Cullen MM, Daddona PE: Conjugation and evaluation of 7E3 × P4B6, a chemically cross-linked bispecific F(ab')₂ antibody which inhibits platelet aggregation and localizes tissue plasminogen activator to the platelet surface. *Bioconj Chem* 3:126, 1992

25. Peterson GL: Determination of total protein. *Methods Enzymol* 91:95, 1983

26. Fraker DJ, Speck JC: Protein and cell membrane iodinations with a sparingly soluble chloroamide. 1,3,4,5-tetrachloro-3a,6a-diphenyl-glycoluril. *Biochem Biophys Res Commun* 80:849, 1978

27. Marguerie GA, Plow EF, Edgington TS: Human platelets possess an inducible and saturable receptor specific for fibrinogen. *J Biol Chem* 254:5357, 1979

28. Lombardo VT, Hodson E, Roberts JP, Kunicki TJ, Zimmerman TS, Ruggeri ZM: Independent modulation of von Willebrand factor and fibrinogen binding to platelet membrane GPIIb/IIIa. *J Clin Invest* 76:1950, 1985

29. Blum M, Jordan R, Wagner C, Mattis J: Immunopurification and functional characterization of platelet GPIIb/IIIa complexes. *Blood* 74:398a, 1989 (abstr no. 1521, suppl 1)

30. Cramer EM, Savidge GF, Vainchenker W, Berndt MC, Pidard D, Caen JP, Masse J-M, Breton-Gorius J: Alpha-granule pool of GPIIb-IIIa in normal and pathologic platelets and megakaryocytes. *Blood* 75:1220, 1990

31. Escolar G, Leistikow E, White JG: The fate of the open canalicular system in surface and suspension-activated platelets. *Blood* 74:1983, 1989

32. Collier BS, Cheresch DA, Asch E, Seligsohn U: Platelet vitronectin receptor expression differentiates Iraqi-Jewish from Arab patients with Glanzmann thrombasthenia in Israel. *Blood* 77:75, 1991

33. Collier BS: Biochemical and electrostatic considerations in primary platelet aggregation. *Ann NY Acad Sci* 416:693, 1983

34. Sosnick TR, Benjamin DC, Novotny J, Seeger PA, Trehwella J: Distances between the antigen-binding sites of three murine antibody subclasses measured using neutron and X-ray scattering. *Biochemistry* 31:1779, 1992

35. Kloczewiak M, Timmons S, Lukas TJ, Hawiger J: Platelets receptor recognition site on human fibrinogen: synthesis and structure-function relationship of peptides corresponding to the carboxyl-terminal segment of the gamma chain. *Biochemistry* 23:1767, 1984

36. Farrell DH, Thiagarajan P, Chung DW, Davie EW: Role of fibrinogen α and γ chain sites in platelet aggregation. *Proc Natl Acad Sci USA* 89:10729, 1992

37. Weisel JW, Nagaswami C, Vilaire G, Bennett JS: Examination of the platelet membrane GPIIb-IIIa complex and its interaction with fibrinogen and other ligands by electron microscopy. *J Biol Chem* 267:16637, 1992



blood[®]

1996 88: 907-914

Analysis of GPIIb/IIIa receptor number by quantification of 7E3 binding to human platelets

CL Wagner, MA Mascelli, DS Neblock, HF Weisman, BS Coller and RE Jordan

Updated information and services can be found at:

<http://www.bloodjournal.org/content/88/3/907.full.html>

Articles on similar topics can be found in the following Blood collections

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>