

HEMOSTASIS, THROMBOSIS, AND VASCULAR BIOLOGY

Glycoprotein Ib-V-IX, a Receptor for von Willebrand Factor, Couples Physically and Functionally to the Fc Receptor γ -Chain, Fyn, and Lyn to Activate Human Platelets

By Shahrokh Falati, Christine E. Edmead, and Alastair W. Poole

The adhesion molecule von Willebrand factor (vWF) activates platelets upon binding 2 surface receptors, glycoprotein (GP) Ib-V-IX and integrin $\alpha_{IIb}\beta_3$. We have used 2 approaches to selectively activate GP Ib using either the snake venom lectin alboaggregin-A or mutant recombinant forms of vWF ($\Delta A1$ -vWF and RGGs-vWF) with selective binding properties to its 2 receptors. We show that activation of GP Ib induces platelet aggregation, secretion of 5-hydroxy tryptamine (5-HT), and an increase in cytosolic calcium. Syk becomes tyrosine phosphorylated and activated downstream of GP Ib, and associates with several tyrosine-phosphorylated proteins including the Fc receptor γ -chain through interaction with Syk SH2 domains. GP Ib physically associates with the γ -chain in GST-Syk-SH2 precipitates from platelets stimulated through GP Ib, and 2 Src family

kinases, Lyn and Fyn, also associate with this signaling complex. In addition, GP Ib stimulation couples to tyrosine phosphorylation of phospholipase C γ 2. The Src family-specific inhibitor PP1 dose-dependently inhibits phosphorylation of Syk, its association with tyrosine-phosphorylated γ -chain, phosphorylation of PLC γ 2, platelet aggregation, and 5-HT release. The results indicate that, upon activation, GP Ib is physically associated with FcR γ -chain and members of the Src family kinases, leading to phosphorylation of the γ -chain, recruitment, and activation of Syk. Phosphorylation of PLC γ 2 also lies downstream of Src kinase activation and may critically couple early signaling events to functional platelet responses.

© 1999 by The American Society of Hematology.

PLATELET ADHESION to subendothelial structures is an early critical event in hemostasis and thrombosis. von Willebrand factor (vWF) is a major adhesive glycoprotein (GP) required for normal hemostasis in conditions of high shear stress,¹⁻⁴ such as occur in small arterioles and arterial capillaries. In the presence of shear stress or modulators such as ristocetin, vWF is able to induce signaling in platelets including hydrolysis of phosphoinositides, a transient increase in cytosolic calcium, activation of protein kinase C, cytoskeletal reorganization, and platelet aggregation.⁵⁻⁹

Platelets have 2 receptors for vWF that are sequentially bound upon interaction with vWF: GP Ib in the GP Ib-V-IX complex and the integrin $\alpha_{IIb}\beta_3$.¹⁰ Previous studies have shown that the tyrosine kinase Syk becomes tyrosine phosphorylated and activated in vWF-stimulated platelets^{11,12} downstream of the primary receptor GP Ib, and Syk has been shown to play an essential role in signaling through another adhesion molecule, collagen.¹³⁻²⁰ Therefore, it was important to establish the mechanism by which Syk is activated downstream of GP Ib in vWF-stimulated platelets. Syk is classically activated by engagement of its tandem Src homology 2 (SH2) domains with doubly phosphorylated tyrosine residues in proteins containing immune-receptor tyrosine-containing activation motifs (ITAMs).²¹ Nei-

ther GP Ib-V-IX nor $\alpha_{IIb}\beta_3$ contain ITAM sequences, and only 2 ITAM-containing proteins have been described in platelets; the Fc receptor γ -chain (FcR γ -chain) and the low-affinity receptor for IgG, Fc γ RIIA.^{13,22,23} It is possible that either or both of these proteins are required for vWF signaling in platelets and there is evidence to show constitutive association of Fc γ RIIA with GP Ib-V-IX.^{24,25} However, the FcR γ -chain has been shown to be critically involved in signaling downstream of collagen¹⁹ and its putative signaling receptor GP VI^{15,16,26} and may, therefore, also be involved in mediating GP Ib signaling to Syk.

We have used 2 approaches to selective activation of GP Ib-V-IX: (1) use of alboaggregin-A, a lectin purified from the venom of the white-lipped tree viper *Trimeresurus albolabris*, which binds to GP Ib inducing activation, and (2) use of mutant recombinant vWFs that bind selectively to either GP Ib or the integrin $\alpha_{IIb}\beta_3$. Recently, a group of viper venom proteins has been reported to interact with GP Ib on platelets resulting in either platelet agglutination and inhibition of ristocetin-induced vWF binding or induction of platelet activation.²⁷⁻³¹ One such protein isolated from the venom of *T albolabris*, the 50-kD C-type lectin alboaggregin-A, has been shown to potentially induce platelet activation through binding to GP Ib.^{28,32} In the present study we have isolated alboaggregin-A to use as a selective tool to induce platelet activation downstream of GP Ib. In addition, we have used 2 recombinant mutant forms of vWF, in combination with the modulator ristocetin, to differentially activate GP Ib or the integrin $\alpha_{IIb}\beta_3$. $\Delta A1$ -vWF is a deletion mutant of wild-type human vWF in which the major part of the A1 domain, responsible for binding GP Ib α , has been ablated (residues 478-716).³³ RGGs-vWF has a point mutation (1746^{D→G}) in the RGD sequence of vWF, disabling its binding to integrin $\alpha_{IIb}\beta_3$,³⁴ while allowing normal binding to GP Ib. Using these 2 approaches, we set out to establish a signaling pathway downstream of GP Ib, investigating the role of Src family kinases, FcR γ -chain, Syk, and PLC γ 2 in functional responses downstream of this receptor.

From the Department of Pharmacology, University of Bristol, School of Medical Sciences, Bristol, UK.

Submitted February 8, 1999; accepted April 26, 1999.

Supported by grants from the Biotechnology and Biological Sciences Research Council, UK and the British Heart Foundation.

Address reprint requests to Alastair W. Poole, PhD, Department of Pharmacology, School of Medical Sciences, University Walk, Bristol BS8 1TD, UK; e-mail: A.Poole@bris.ac.uk.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

© 1999 by The American Society of Hematology.

0006-4971/99/9405-0029\$3.00/0

MATERIALS AND METHODS

Materials. Plasma vWF (pvWF) and 2 mutant recombinant forms of vWF ($\Delta A1$ -vWF and RGGG-vWF) were kind gifts from Prof J.J. Sixma and Dr T. Vink (Utrecht, The Netherlands), and were prepared as previously described.^{33,34} *T albolabris* venom was a generous gift from Prof R.D.G. Theakston (Liverpool, UK). The anti-GP Ib mouse monoclonal antibody, MoAb 6D1, was a kind gift from Prof B. Collier (New York, NY). Antiphosphotyrosine MoAb 4G10 and polyclonal anti-FcR γ -chain antibody were from Upstate Biotechnology Inc (TCS Biologicals Ltd, Bucks, UK). Polyclonal anti-Syk, anti-Lyn, anti-GP Ib, anti-Fyn, and anti-PLC γ 2 antibodies were from Santa Cruz Biotechnology (Autogen Bioclear, Calne, Wiltshire, UK). GST-fusion protein of Syk tandem SH2 domains (GST-Syk-SH2) was kindly supplied by Dr S. Watson (Oxford, UK). A suspension of type I collagen fibers from equine tendon (Horm collagen) was from Nycomed (Munich, Germany). γ -[³²P]-ATP and enhanced chemiluminescence (ECL) reagents were from Amersham Plc (Amersham, UK). Ristocetin, protein A-Sepharose CL 4B, Tween 20, and phenylmethylsulfonyl fluoride (PMSF) were from Sigma (Poole, Dorset, UK). Acrylamide/bisacrylamide solution was from National Diagnostics (Hull, UK). The Src family kinase inhibitor PP1 was from Alexis Corp (Nottingham, UK). All other reagents were of analytical grade.

Preparation and stimulation of human platelets. Human blood was drawn from drug-free volunteers on the day of the experiment using acid citrate dextrose (ACD: 120 mmol/L sodium citrate, 110 mmol/L glucose, 80 mmol/L citric acid 1:7, vol/vol) as anticoagulant. Platelet-rich plasma (PRP) was prepared by centrifugation (200g, 20 minutes) and platelets were isolated by centrifugation of PRP (1,000g, 10 minutes) in the presence of prostacyclin (0.1 μ g/mL). The pellet was resuspended to a density of 4.10^8 platelets/mL in modified Tyrode's-HEPES buffer (145 mmol/L NaCl, 2.9 mmol/L KCl, 10 mmol/L HEPES, 1 mmol/L MgCl₂, 5 mmol/L glucose, pH 7.3). Indomethacin (10 μ mol/L) was added to platelet suspensions throughout subsequent procedures. Stimulation of platelets was performed at 37°C in an aggregometer with continuous stirring at 800 rpm. Concentrations of pvWF (10 μ g/mL) and RGGG-vWF (6 μ g/mL) were shown to give near-maximal responses, while $\Delta A1$ -vWF was used at 3 μ g/mL because it had been shown that this concentration allows maximal binding to integrin $\alpha_{IIb}\beta_3$.³³

Platelet aggregation and release of 5-HT. Platelet aggregation was measured by optical turbidometry³⁵ using a platelet aggregometer (Chronolog Corp, Havertown, PA). For aggregation studies, platelets were suspended in Tyrode's-HEPES without EGTA, with the exception of those experiments conducted to distinguish fibrinogen-dependent aggregation from adhesion, where platelets were preincubated with EGTA (1 mmol/L) to block fibrinogen binding. The data shown represent the decrease in optical density as a percentage of the maximum possible decrease. For assessment of 5-HT release, platelets were loaded with [³H]-5-HT by incubation with 0.2 μ Ci/mL PRP for 1 hour at 37°C. Platelets were preincubated with EGTA (1 mmol/L) before stimulation to prevent aggregation, and the reaction was terminated by brief microcentrifugation and [³H]-5-HT release into supernatant was determined by scintillation spectrometry. [³H]-5-HT release was expressed as a percentage of the total tissue content as described previously.¹⁹

Measurement of cytosolic [Ca²⁺]. This was performed as previously described.³⁶ Briefly, PRP was incubated with Fura-2-AM (3 μ mol/L) at 30°C for 45 minutes, and platelets prepared as described above. Platelets were stimulated with various concentrations of alboaggregin-A under stirred conditions at room temperature in the absence of EGTA. Fluorescence excitation was made at wavelengths 340 and 380 nm, and emission at 510 nm was measured using a Perkin-Elmer LS50B spectrofluorimeter. Data are presented as the excitation fluorescence ratio (340:380 nm).

Purification of alboaggregin-A. Alboaggregin-A was prepared from the crude venom of the viper *T albolabris*, as previously described^{28-30,32} by ion exchange and phenyl sepharose hydrophobic chromatography, and showed physical and functional characteristics identical to that purified by other workers.^{28-30,32} Briefly, the protein had an apparent molecular weight of 50 kD by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) under nonreduced conditions and induced platelet aggregation and 5-HT release in a manner not blocked by the thrombin protease inhibitor PPACK (data not shown). Functional characteristics of alboaggregin-A are described in Results.

Immunoblotting. Platelets were activated in the presence of EGTA (1 mmol/L) for all blotting and protein precipitation studies. Reactions were stopped by adding an equal volume of Laemmli buffer (2X) and samples were heated for 5 minutes at 95°C. Proteins were separated by either 10% SDS-PAGE or by SDS-PAGE on 10% to 18% gradient slab gels and transferred to polyvinylidene difluoride (PVDF) blotting membranes using a semi-dry transfer system (60 minutes, 15 V). Membranes were incubated for 60 minutes at room temperature with primary followed by secondary antibodies and detected by ECL (Amersham, UK).

Immunoprecipitation and GST-fusion protein precipitation. Reactions were stopped by lysis with an equal volume of 2X extraction buffer (2% [vol/vol] Triton X-100, 300 mmol/L NaCl, 20 mmol/L Tris, 1 mmol/L phenylmethylsulfonyl fluoride [PMSF], 10 mmol/L EDTA, 2 mmol/L Na₃VO₄, 10 μ g/mL leupeptin, 10 μ g/mL aprotinin, 1 μ g/mL pepstatin A, pH 7.3), and insoluble material removed by centrifugation (13,000g, 5 minutes, 4°C). Supernatants were then precleared by incubation with protein A-Sepharose (PAS) for immunoprecipitation experiments, or with glutathione-agarose for GST-fusion protein precipitations, for 1 hour at 4°C, followed by centrifugation (13,000g, 5 minutes, 4°C). For immunoprecipitations, supernatants were then incubated with PAS and the appropriate immunoprecipitating antibody for 120 minutes at 4°C, while for GST fusion protein precipitations, supernatants were incubated with glutathione-agarose beads and GST-Syk-SH2 (10 μ g/mL) for 120 minutes at 4°C. Beads were then washed once in extraction buffer and twice more in TBS-T before addition of Laemmli sample-treatment buffer. Precipitated proteins were then subjected to SDS-PAGE, transferred to PVDF membrane, and probed with appropriate antibodies as described in Immunoblotting.

In vitro kinase assay. Immunoprecipitated Syk was suspended in 20 μ L kinase assay buffer (5 mmol/L MgCl₂, 5 mmol/L MnCl₂, 100 mmol/L NaCl, 10 μ mol/L adenosine triphosphate (ATP), 20 mmol/L HEPES at pH 7.2) and the reaction started by addition of [γ -³²P]-ATP (250 μ Ci/mL). After incubation for 15 minutes at 25°C, the reaction was terminated by addition of 0.5 mL ice-cold 100 mmol/L EDTA. Samples were subjected to SDS-PAGE and phosphorylated proteins were visualized by autoradiography.

RESULTS

Alboaggregin-A induces platelet aggregation, release of 5-HT, and an increase in cytosolic calcium through activation of GP Ib. A concentration-response relationship was determined for induction of secretion of 5-HT (Fig 1A) by alboaggregin-A (60 seconds), and the EC₅₀ concentration of 3.5 μ g/mL was used throughout the rest of the study to activate platelets, unless otherwise stated. At this concentration, alboaggregin-A induced platelet aggregation, which was substantially inhibited by calcium chelation with EGTA (1 mmol/L) or the monoclonal anti-GP Ib blocking antibody 6D1 (9 μ g/mL) (Fig 1B) demonstrating platelet activation through binding GP Ib. In contrast, aggregation in response to collagen (100 μ g/mL, 2 minutes) was not inhibited in the presence of 6D1 (10 μ g/mL). EGTA




Fig 1. Alboaggregin-A induces platelet aggregation, release of 5-HT and an increase in cytosolic calcium upon binding to GP Ib. For 5-HT release, studies platelets were preloaded with [^3H]-5-HT and stimulated with various concentrations of alboaggregin-A (A, D) or 3.5 $\mu\text{g/ml}$ alboaggregin-A (B, C) for 1 minute. Collagen (100 $\mu\text{g/ml}$; 2 minutes) was used for comparison and as a negative control (B, C). Release of 5-HT is presented as a percentage of total 5-HT content, and a dose-response relationship was determined in (A), showing an EC_{50} of 3.5 $\mu\text{g/ml}$. Platelet aggregation is presented as the decrease in optical density induced by agonist as a percentage of the maximal possible decrease (B). Platelets were pretreated with MoAb 6D1 (9 $\mu\text{g/ml}$) for 6 minutes, or EGTA (1 mmol/L) for 10 minutes, before stimulation with agonist (B, C). For assessment of changes in cytosolic calcium (D), platelets were preloaded with the calcium indicator dye Fura-2 and fluorescence measurements were made at emission wavelength 510 nm. Data are presented as the ratio of fluorescence measurements at excitation wavelengths 340 and 380 nm. Alboaggregin-A is added at the time point indicated by the arrow, at concentrations indicated on the right of the graphs. Data presented are means \pm SEM for 3 experiments (A, B, C) or are representative of 3 separate experiments (D).

(1 mmol/L) partially inhibited platelet aggregation by collagen, the remaining response being caused by calcium-independent adhesion of platelets to collagen fibers. Secretion of 5-HT induced by alboaggregin-A was inhibited by MoAb 6D1 (Fig 1C). In contrast with aggregation, however, secretion was unaffected by EGTA, demonstrating alboaggregin-A to bind GP Ib in the absence of extracellular calcium. However, collagen-induced secretion of 5-HT was unaffected by either EGTA or MoAb 6D1. Alboaggregin-A also induced a dose-dependent rapid increase in cytosolic calcium, as assessed by the change in 340:380-nm fluorescence ratio of Fura-2-loaded platelets (Fig 1D).

Alboaggregin-A induces tyrosine phosphorylation of Syk and PLC γ 2 and activation of Syk. Previous reports have shown that a variety of platelet agonists, including collagen, thrombin, and activation of GP Ib, are able to induce the tyrosine phosphorylation and activation of Syk.^{11-13,19,37} Figure 2A shows that both alboaggregin-A and collagen, which was used for comparison, induced tyrosine phosphorylation of Syk above

basal levels. In addition, Syk activity, assayed *in vitro* as autophosphorylation, was shown to increase markedly upon activation with alboaggregin-A (Fig 2B). Tyrosine phosphorylation of PLC γ 2 has previously been reported in response to collagen and crosslinking Fc γ RIIA in platelets,^{13,14,20,38,39} and here we show that stimulation of platelets with alboaggregin-A caused tyrosine phosphorylation of PLC γ 2 (Fig 2C). Collagen-induced phosphorylation of PLC γ 2 is shown for comparison. Alboaggregin-A induced greater phosphorylation of Syk than collagen, possibly due to phosphorylation of additional tyrosine residues not phosphorylated upon collagen stimulation.

Syk associates with FcR γ -chain, mediated by tandem SH2 domains of Syk. Figure 3A shows that alboaggregin-A induced a marked, early association between FcR γ -chain and GST-Syk-SH2, which was likely to be caused by a rapid phosphorylation of FcR γ -chain. The association decreased at later time points, probably reflecting a time-dependent dephosphorylation of the γ -chain. In addition, Fig 3B shows that pVWF (10 $\mu\text{g/ml}$) in the presence of ristocetin (1 mg/ml) stimulated a




Fig 2. Alboaggregin-A induces tyrosine phosphorylation of Syk and PLC γ 2 and activates Syk kinase. For (A), Syk and for (C), PLC γ 2 were immunoprecipitated from lysates of basal platelets (lane 1) or platelets stimulated with either alboaggregin-A (3.5 μ g/mL; lane 2) for 1 minute or collagen (100 μ g/mL; lane 3) for 2 minutes and Western blotted with either 4G10 [A(i) and C(i)] and anti-Syk [A(ii)] or anti-PLC γ 2 [C(ii)]. For (B), a kinase assay was performed in vitro on Syk immunoprecipitates from basal platelets (lane 1) or platelets stimulated with alboaggregin-A (3.5 μ g/mL; lane 2) for 1 minute, and presented as an autoradiograph. The results are representative of 3 separate experiments.

marked, early association of tyrosine phosphorylated γ -chain with GST-Syk-SH2, which diminished with time. Additional tyrosine-phosphorylated proteins of 44, 56, and 59 kDa were found to associate with GST-Syk-SH2 both in alboaggregin-A-stimulated and vWF-stimulated platelets, although these proteins remain unidentified. This finding leaves the possibility that there are routes other than the γ -chain by which Syk may be activated downstream of GP Ib. Precipitates from collagen-stimulated platelets are shown for comparison.

vWF induces Syk phosphorylation and association with FcR γ -chain through activation of GP Ib. To show that vWF-stimulated association was downstream of GP Ib, Syk was immunoprecipitated from platelets challenged with mutant vWFs that differentially bind either GP Ib or the integrin $\alpha_{IIb}\beta_3$. Figure 4A shows that although pvWF induced Syk phosphorylation and association of Syk with tyrosine-phosphorylated γ -chain, this response was absent in platelets activated by Δ A1-vWF, a mutant unable to bind GP Ib, in the presence of ristocetin (1 mg/mL). On the other hand, RGGs-vWF, a mutant that is unable to bind $\alpha_{IIb}\beta_3$ but is able to bind GP Ib, induced




Fig 3. Alboaggregin-A and pvWF induce association of tyrosine phosphorylated FcR γ -chain with GST-Syk-SH2. Platelets were stimulated with either alboaggregin-A (3.5 μ g/mL), pvWF (10 μ g/mL), and ristocetin (1 mg/mL) or with collagen (100 μ g/mL) for the indicated times, and proteins were precipitated from cell lysates using 10 μ g of GST-Syk SH2 per lane. Precipitated proteins were separated by SDS-PAGE and immunoblotted with 4G10 [A(i) and B(i)] and anti-FcR γ -chain [A(ii) and B(ii)]. For (A), lane 1 is resting platelets and lanes 2 through 4 were stimulated with alboaggregin-A for the times indicated. For (B), lanes 1 and 7 were resting platelets, lanes 2 through 6 were stimulated with pvWF in the presence of ristocetin for the times indicated, and in lane 8, platelets were stimulated with collagen. Results shown are representative of at least 3 separate experiments.




Fig 4. Mutant forms of vWF induce differential association between Syk and tyrosine phosphorylated FcR γ -chain. Platelets were stimulated with mutant vWF or pvWF in the presence of ristocetin for 45 seconds, thrombin for 45 seconds, or collagen for 120 seconds. Syk was immunoprecipitated, run on SDS-PAGE, and Western blotted with 4G10 [A(i) and B(i)] or with anti-Syk [A(ii) and B(ii)]. [A(i)]: lane 1, resting platelets; lane 2, pvWF (10 μ g/mL) and ristocetin (1 mg/mL); lane 3, Δ A1-vWF (3 μ g/mL) and ristocetin (1 mg/mL). [B(i)] Phosphorylation of FcR γ -chain which has been precipitated in association with Syk from basal platelets (lane 1); platelets stimulated with pvWF (10 μ g/mL, lane 2), Δ A1-vWF (3 μ g/mL, lane 3), RGGG-vWF (6 μ g/mL, lane 4) in the presence of ristocetin (1 mg/mL); thrombin (1 U/mL, lane 5) or collagen (100 μ g/mL, lane 6). Equal amounts of Syk were present in each lane [B(ii)]. Immunoblots shown are representative of 4 separate experiments.

both tyrosine phosphorylation of Syk (data not shown) and association with the γ -chain (Fig 4B). Furthermore, as has previously been reported,^{13,14,16,20,37} both thrombin and collagen were able to induce tyrosine phosphorylation of Syk (data not shown). However, Syk associated with FcR γ -chain only in collagen-stimulated, but not thrombin-stimulated, platelets (Fig 4B). These findings indicate that FcR γ -chain associates with

Syk only in cells stimulated through GP Ib or by collagen, but not by thrombin.

FcR γ -chain associates with Fyn, Lyn, and GP Ib in platelets activated with alboaggregin-A. Recent studies have reported a selective role for Fyn and Lyn, but not other Src-family kinase members, in platelets stimulated through the collagen receptor GP VI.^{14,40} We used GST-Syk-SH2 to precipitate γ -chain from alboaggregin-A-activated platelets, and identified co-associating proteins by Western blotting. Under these conditions, both Fyn and Lyn associated with GST-Syk-SH2 complexed to FcR γ -chain (Fig 5); in addition, GP Ib precipitated with this signaling complex, showing a physical association between these signaling elements under stimulated conditions.

PP1 inhibits Syk phosphorylation, association with tyrosine-phosphorylated γ -chain, and phosphorylation of PLC γ 2. The Src family kinase inhibitor PP1 was shown to dose-dependently inhibit phosphorylation of multiple proteins induced by alboaggregin-A (data not shown). Figure 6A shows that PP1 inhibited phosphorylation of Syk and its association with tyrosine-phosphorylated γ -chain in a concentration-dependent manner. Interestingly, 5 μ mol/L PP1 almost abolished γ -chain phosphorylation and association, but only partially inhibited Syk phosphorylation. This may provide further evidence for the existence of γ -chain-independent pathways to activation of Syk by GP Ib.

Figure 6B shows that phosphorylation of PLC γ 2 was dose-dependently inhibited by PP1 such that there was partial inhibition at 5 μ mol/L PP1 and full inhibition by 20 μ mol/L PP1. This result parallels that for Syk phosphorylation and is consistent with PLC γ 2 activation being closely coupled to Syk activation.

Inhibition of Src family kinases inhibits GP Ib-mediated platelet aggregation and secretion of 5-HT. To demonstrate a functional requirement for Src family kinases in platelet activation through GP Ib, we showed that PP1 dose-dependently




Fig 5. Coprecipitation of FcR γ -chain, GPIb α , Lyn, and Fyn with GST-Syk-SH2 from platelets stimulated with alboaggregin-A. Platelets were either unstimulated (lane 1) or were stimulated with alboaggregin-A (3.5 μ g/mL) for 15 seconds (lane 2) or 60 seconds (lane 3), and proteins were precipitated with 10 μ g of GST-Syk-SH2 per lane. Precipitated proteins were then separated by SDS-PAGE and immunoblotted with α GPIb α , α Lyn, α Fyn, or α FcR γ -chain, as indicated. Results shown are representative of 3 separate experiments.




Fig 6. PP1 dose-dependently inhibits alboaggregin-A-induced tyrosine phosphorylation of FcR γ -chain, Syk, and PLC γ 2. Platelets were preincubated for 3 minutes at 37°C with 0.25% dimethyl sulfoxide (DMSO) or various concentrations of PP1 before stimulation with alboaggregin-A (3.5 μ g/mL) for 1 minute. Platelet suspensions were then lysed and either Syk or PLC γ 2 immunoprecipitated. Precipitated proteins were then separated by SDS-PAGE and immunoblotted with 4G10 [A(i)] and B(i)] and subsequently with either anti-Syk [A(ii)] or anti-PLC γ 2 [B(ii)]. Immunoblots shown are representative of 3 separate experiments.

inhibited both platelet aggregation and release of 5-HT induced by alboaggregin-A (Fig 7). Full inhibition of responses was achieved by 20 μ mol/L PP1, in parallel with inhibition of tyrosine phosphorylation of Syk and PLC γ 2.

DISCUSSION

It has become clear that vWF is able to induce platelet activation,^{5,6,41,42} and it is now emerging that tyrosine phosphorylation events may play a central signaling role in vWF-induced activation.^{1,43-45} Furthermore, many of these signaling events are directly downstream of the primary vWF receptor GP Ib-V-IX, which has been shown to induce tyrosine phosphorylation of multiple platelet proteins.⁴³⁻⁴⁵ Recent evidence suggests that vWF may activate the nonreceptor tyrosine kinase Syk through GP Ib. Antibody-induced cross-linking of GP Ib induces a small aggregation response associated with activation of Syk¹² and, in the presence of the modulator botrocetin, vWF

induces tyrosine phosphorylation and activation of Syk downstream of GP Ib.¹¹

The aim of this report was to elucidate mechanisms that couple GP Ib to Syk activation, and to determine signaling events downstream of Syk that may regulate functional activities. Although there may be other mechanisms by which Syk may be activated,^{46,47} it is generally recognized that Syk forms signaling complexes at the plasma membrane by interaction of its tandem Src homology 2 (SH2) domains with proteins containing phosphorylated ITAMs.^{21,48,49} None of the components of GP Ib-V-IX complex possess ITAM motifs; however, platelets express at least 2 ITAM-containing proteins; the low-affinity receptor for IgG, Fc γ RIIA, and FcR γ -chain.^{13,22,23} It is possible that GP Ib-V-IX forms a functional complex with either or both of these proteins, allowing it to couple to Syk upon activation.

The FcR γ -chain forms an integral part of several antibody Fc receptors and has recently been shown to be complexed constitutively to the platelet receptor for the adhesion molecule collagen, GP VI.^{15,26} It becomes tyrosine phosphorylated and associates with Syk upon activation of platelets with collagen or the GP VI-specific C-type lectin convulxin.^{15,16,23,26,50} In knockout studies, both the γ -chain and Syk have been shown to be essential for activation of platelets by collagen¹⁹ and, importantly in the absence of FcR γ -chain, collagen is unable to induce Syk activation, showing the FcR γ -chain to be an essential upstream regulator of Syk. We were therefore interested in investigating whether Syk activation downstream of GP Ib might involve an essential receptor complex with the FcR γ -chain.

Several tyrosine-phosphorylated proteins associate with Syk upon activation with vWF or alboaggregin-A, and prominent amongst these is a doublet of 14 kD that was shown to be the FcR γ -chain. The association was through the SH2 domains of




Fig 7. PP1 dose-dependently inhibits alboaggregin-A-induced platelet aggregation and 5-HT secretion. Platelets were preincubated for 3 minutes at 37°C with 0.25% DMSO or various concentrations of PP1 (indicated on the right of the graphs) before stimulation with alboaggregin-A (3.5 μ g/mL) for 1 minute. Platelet aggregation was measured as a decrease in optical density of a stirred platelet suspension and 5-HT release was measured after loading cells with [³H]-5-HT. Both platelet aggregation and release of 5-HT are presented as percentages of the responses induced in the absence of PP1. Data presented are means \pm SEM for 3 experiments.

Syk and was absent in experiments using the $\Delta A1$ -vWF mutant, which is unable to bind GP Ib, but maintained with RGGG-vWF, which is unable to bind the integrin $\alpha_{IIb}\beta_3$, thus strongly supporting the hypothesis that the FcR γ -chain couples GP Ib to Syk. Interestingly, although thrombin was also able to induce phosphorylation of Syk, in agreement with other investigators,^{47,51} there was no association with the γ -chain, indicating a γ -chain-independent mechanism to Syk activation downstream of this receptor. Collagen, on the other hand, induced phosphorylation of Syk that associated with phosphorylated FcR γ -chain, in agreement with previous reports.^{14,19,23} Therefore, it is clear that FcR γ -chain is likely to represent a major pathway to activation of Syk downstream of GP Ib, but the presence of other tyrosine-phosphorylated proteins precipitated by Syk SH2 domains leaves open the possibility that there are other routes to Syk activation independent of FcR γ -chain, downstream of GP Ib. This would be consistent with phenotypes in knockout mice that we observed previously.¹⁹ In the absence of Syk, mice suffer a severe petechial hemorrhaging during fetal development^{19,52,53} and platelets from these mice do not respond to collagen. In contrast, although platelets from mice lacking FcR γ -chain also lack a response to collagen, they do not manifest a major bleeding disorder, suggesting multiple mechanisms by which Syk may be activated downstream of several receptors. One possible link to Syk in human platelets would be Fc γ RIIA, which has been shown to be constitutively associated with GP Ib,^{24,25} although this would not explain the difference in knockout phenotypes because mice do not possess the Fc γ RIIA gene.⁵⁴

It is now established that Src family kinases are essential for primary signals to Syk through ITAM motif-containing proteins.^{21,55,56} Platelets express abundant quantities of several Src family kinases including Src, Fyn, Lyn, Yes, Hck, Fgr, and Lck.⁵⁷⁻⁶⁰ It has previously been shown that Src transiently associates with Syk in vWF-activated platelets,¹¹ and that Src may be activated by vWF, translocating to the cytoskeleton upon activation.⁶¹ In the present study we show that both Fyn and Lyn physically associate with GST-Syk-SH2:FcR γ -chain complex upon platelet activation through GP Ib. In addition, GP Ib itself forms a component of this signaling complex. The technique used to precipitate FcR γ -chain in these studies, in which we use GST-Syk-SH2 to precipitate tyrosine-phosphorylated FcR γ -chain, only allows precipitation of γ -chain from activated cells. This limitation does not allow us to conclude whether the association between FcR γ -chain and the other signaling components takes place constitutively or upon platelet activation. However, there is recent evidence that Src-family kinases may associate with nonphosphorylated γ -chain,⁶² and we present evidence that Fyn may bind to GST-Syk-SH2 constitutively (see Fig 5). We therefore speculate that Fyn may form a physical link between the γ -chain and Syk under basal conditions. We also have preliminary evidence that FcR γ -chain is constitutively associated to GP Ib because it is isolated from unstimulated cell lysates precipitated with alboaggregin-A bound to a solid phase (S.F., A.W.P., unpublished observations, September 1998). Taken together, these results may parallel previous data for the platelet collagen receptor, GP VI, in which FcR γ -chain associates with the receptor and Src family members Fyn and Lyn irrespective of activation.¹⁴

Therefore, it is concluded that GP Ib, γ -chain, Fyn, Lyn, and Syk form a physical complex upon activation of GP Ib, leading to multiple functional events in platelets including platelet aggregation, 5-HT release, and an increase in cytosolic calcium. We were interested to establish whether the receptor-signaling complex would also signal to activation of PLC γ 2, which has previously been shown to be tyrosine phosphorylated in collagen-stimulated platelets.^{13,38,39} PLC γ 2 has been shown to be downstream of Syk in hematopoietic cells including platelets,^{19,63-65} and may functionally couple the proximal GP Ib signaling complex to downstream functional events. In this report we show that PLC γ 2 becomes tyrosine phosphorylated downstream of GP Ib. In addition, Src family kinases are essential for its tyrosine phosphorylation because PP1, which has selectivity for Src family kinases over Syk at concentrations up to 100 μ mol/L,^{55,66} dose-dependently inhibits PLC γ 2 phosphorylation as well as Syk phosphorylation, platelet aggregation, and secretion of 5-HT. FcR γ -chain phosphorylation is also inhibited by PP1, although at 5 μ mol/L the inhibition of γ -chain phosphorylation is greater than that for Syk or PLC γ 2. This disproportion between the phosphorylation of these proteins may provide further evidence that Syk is activated by γ -chain-independent pathways, such as that described by Gao et al,⁴⁶ or there may be sufficient signal amplification at the level of the γ -chain. It is also interesting that 20 μ mol/L PP1, which blocks phosphorylation of all signaling proteins studied here, does not fully block 5-HT release. This provides evidence for Src-family kinase-independent pathways to platelet activation by GP Ib. These pathways may include PI 3-kinase, which has been shown to be activated by vWF,⁶¹ leading to direct activation of PLC γ 2⁶⁷⁻⁷¹ without a requirement for tyrosine phosphorylation. 14-3-3 proteins have also been shown to bind GP Ib-V-IX components⁷²⁻⁷⁵ and may form an important Src-family kinase-independent signaling pathway.

Based on the present findings, we put forward a working model in which GP Ib associates with FcR γ -chain, either constitutively or upon activation of the receptor, and that Src family kinases Fyn and Lyn phosphorylate FcR γ -chain leading to binding of Syk through its tandem SH2 domains, activation of Syk, and finally PLC γ 2 tyrosine phosphorylation and activation leading to platelet functional responses. However, this model leaves several important questions to be addressed; the possibility of FcR γ -chain-independent mechanisms by which GP Ib can couple to Syk activation warrants further investigation. Furthermore, details of the construction of the receptor-FcR γ -chain-Fyn-Lyn-Syk complex, under basal and stimulated conditions, remain to be elucidated. Recent studies have also established that PLC γ 2 can become activated in tyrosine phosphorylation-dependent and -independent manners.⁶⁷⁻⁷¹ It remains unknown whether PLC γ 2 becomes activated downstream of GP Ib and, if so, whether tyrosine phosphorylation of PLC γ 2 is the sole mechanism by which PLC γ 2 may become activated downstream of this receptor.

ACKNOWLEDGMENT

The authors are grateful to Prof Jan Sixma and Dr Tom Vink for the kind gift of pvWF and mutant recombinant forms of vWF; Prof David Theakston for generously supplying venom from the viper *Talbolabris*; Dr Jon Gibbins for his time, resources, and expertise in assisting

preparation of purified alboaggregin-A; Prof Barry Collier for kindly supplying MoAb 6D1; and Dr Steve Watson for the generous donation of GST-Syk-SH2.

REFERENCES

- Ruggeri ZM: Mechanisms initiating platelet thrombus formation. *Thromb Haemost* 78:611, 1997
- Ruggeri ZM, Ware J: von Willebrand factor. *FASEB J* 7:308, 1993
- Fressinaud E, Meyer D: von Willebrand factor and platelet interactions with the vessel wall. *Blood Coagul Fibrinol* 2:333, 1991
- Ikeda Y, Handa M, Kawano K, Kamata T, Murata M, Araki Y, Anbo H, Kawai Y, Watanabe K, Itagaki I, Sakai K, Ruggeri ZM: The role of von Willebrand factor and fibrinogen in platelet aggregation under varying shear stress. *J Clin Invest* 87:1234, 1991
- Kroll MH, Harris TS, Moake JL, Handin RI, Schafer AI: von Willebrand factor binding to platelet GPIb initiates signals for platelet activation. *J Clin Invest* 88:1568, 1991
- Chow TW, Hellums JD, Moake JL, Kroll MH: Shear stress-induced von Willebrand factor binding to platelet glycoprotein Ib initiates calcium influx associated with aggregation. *Blood* 80:113, 1992
- Kroll MH, Hellums JD, Guo Z, Durante W, Razdan K, Hrbolich JK, Schafer AI: Protein kinase C is activated in platelets subjected to pathological shear stress. *J Biol Chem* 268:3520, 1993
- Cunningham JG, Meyer SC, Fox JE: The cytoplasmic domain of the α -subunit of glycoprotein (GP) Ib mediates attachment of the entire GP Ib-IX complex to the cytoskeleton and regulates von Willebrand factor-induced changes in cell morphology. *J Biol Chem* 271:11581, 1996
- Yuan Y, Dopheide SM, Ivanidis C, Salem HH, Jackson SP: Calpain regulation of cytoskeletal signaling complexes in von Willebrand factor-stimulated platelets. Distinct roles for glycoprotein Ib-V-IX and glycoprotein IIb-IIIa (integrin $\alpha_{IIb}\beta_3$) in von Willebrand factor-induced signal transduction. *J Biol Chem* 272:21847, 1997
- Savage B, Shattil SJ, Ruggeri ZM: Modulation of platelet function through adhesion receptors. A dual role for glycoprotein IIb-IIIa (integrin $\alpha_{IIb}\beta_3$) mediated by fibrinogen and glycoprotein Ib-von Willebrand factor. *J Biol Chem* 267:11300, 1992
- Asazuma N, Ozaki Y, Satoh K, Yatomi Y, Handa M, Fujimura Y, Miura S, Kume S: Glycoprotein Ib-von Willebrand factor interactions activate tyrosine kinases in human platelets. *Blood* 90:4789, 1997
- Yanabu M, Ozaki Y, Nomura S, Miyake T, Miyazaki Y, Kagawa H, Yamanaka Y, Asazuma N, Satoh K, Kume S, Komiyama Y, Fukuhara S: Tyrosine phosphorylation and p72syk activation by an anti-glycoprotein Ib monoclonal antibody. *Blood* 89:1590, 1997
- Asselin J, Gibbins JM, Achison M, Lee YH, Morton LF, Farndale RW, Barnes MJ, Watson SP: A collagen-like peptide stimulates tyrosine phosphorylation of Syk and phospholipase C γ 2 in platelets independent of the integrin $\alpha_2\beta_1$. *Blood* 89:1235, 1997
- Ezumi Y, Shindoh K, Tsuji M, Takayama H: Physical and functional association of the Src family kinases Fyn and Lyn with the collagen receptor glycoprotein VI-Fc receptor gamma chain complex on human platelets. *J Exp Med* 188:267, 1998
- Gibbins JM, Okuma M, Farndale R, Barnes M, Watson SP: Glycoprotein VI is the collagen receptor in platelets which underlies tyrosine phosphorylation of the Fc receptor γ -chain. *FEBS Lett* 413:255, 1997
- Ichinohe T, Takayama H, Ezumi Y, Arai M, Yamamoto N, Takahashi H, Okuma M: Collagen-stimulated activation of Syk but not c-Src is severely compromised in human platelets lacking membrane glycoprotein VI. *J Biol Chem* 272:63, 1997
- Kralisz U, Cierniewski CS: Tyrosine phosphorylation events during different stages of collagen-induced platelet activation. *Biochim Biophys Acta* 1405:128, 1998
- Polgar J, Clemetson JM, Kehrel BE, Wiedemann M, Magnenat EM, Wells TNC, Clemetson KJ: Platelet activation and signal transduction by convulxin, a C-type lectin from *Crotalus durissus terrificus* (tropical rattlesnake) venom via the p62/GPVI collagen receptor. *J Biol Chem* 272:13576, 1997
- Poole A, Gibbins JM, Turner M, van Vugt MJ, van de Winkel JG, Saito T, Tybulewicz VL, Watson SP: The Fc receptor γ -chain and the tyrosine kinase Syk are essential for activation of mouse platelets by collagen. *EMBO J* 16:2333, 1997
- Yanaga F, Poole A, Asselin J, Blake R, Schieven GL, Clark EA, Law CL, Watson SP: Syk interacts with tyrosine-phosphorylated proteins in human platelets activated by collagen and cross-linking of the Fc γ -IIA receptor. *Biochem J* 311:471, 1995
- Cambier JC: Antigen and Fc receptor signaling. The awesome power of the immunoreceptor tyrosine-based activation motif (ITAM). *J Immunol* 155:3281, 1995
- Chacko GW, Brandt JT, Coggeshall KM, Anderson CL: Phosphoinositide 3-kinase and p72(syk) noncovalently associate with the low affinity Fc γ receptor on human platelets through an immunoreceptor tyrosine-based activation motif: Reconstitution with synthetic phosphopeptides. *J Biol Chem* 271:10775, 1996
- Gibbins J, Asselin J, Farndale R, Barnes M, Law CL, Watson SP: Tyrosine phosphorylation of the Fc receptor γ -chain in collagen-stimulated platelets. *J Biol Chem* 271:18095, 1996
- Moore A, Ross GD, Nachman RL: Interaction of platelet membrane receptors with von Willebrand factor, ristocetin, and the Fc region of immunoglobulin G. *J Clin Invest* 62:1053, 1978
- Sullam PM, Hyun WC, Szollosi J, Dong J, Foss WM, Lopez JA: Physical proximity and functional interplay of the glycoprotein Ib-IX-V complex and the Fc receptor Fc γ RIIA on the platelet plasma membrane. *J Biol Chem* 273:5331, 1998
- Tsuji M, Ezumi Y, Arai M, Takayama H: A novel association of Fc receptor γ -chain with glycoprotein VI and their co-expression as a collagen receptor in human platelets. *J Biol Chem* 272:23528, 1997
- Clemetson KJ, Polgar J, Clemetson JM: Snake venom C-type lectins as tools in platelet research. *Platelets* 9:165, 1998
- Kowalska MA, Tan L, Holt JC, Peng M, Karczewski J, Calvete JJ, Niewiarowski S: Alboaggregins A and B. Structure and interaction with human platelets. *Thromb Haemost* 79:609, 1998
- Peng M, Lu W, Kirby EP: Alboaggregin-B: A new platelet agonist that binds to platelet membrane glycoprotein Ib. *Biochemistry* 30:11529, 1991
- Peng M, Lu W, Kirby EP: Characterization of three alboaggregins purified from *Trimeresurus albolabris* venom. *Thromb Haemost* 67:702, 1992
- Fujimura Y, Kawasaki T, Titani K: Snake venom proteins modulating the interaction between von Willebrand factor and platelet glycoprotein Ib. *Thromb Haemost* 76:633, 1996
- Andrews RK, Kroll MH, Ward CM, Rose JW, Scarborough RM, Smith AI, Lopez JA, Berndt MC: Binding of a novel 50-kilodalton alboaggregin from *Trimeresurus albolabris* and related viper venom proteins to the platelet membrane glycoprotein Ib-IX-V complex. Effect on platelet aggregation and glycoprotein Ib-mediated platelet activation. *Biochemistry* 35:12629, 1996
- Sixma JJ, Schiphorst ME, Verweij CL, Pannekoek H: Effect of deletion of the A1 domain of von Willebrand factor on its binding to heparin, collagen and platelets in the presence of ristocetin. *Eur J Biochem* 196:369, 1991
- Lankhof H, Wu YP, Vink T, Schiphorst ME, Zerwes HG, de Groot PG, Sixma JJ: Role of the glycoprotein Ib-binding A1 repeat and the RGD sequence in platelet adhesion to human recombinant von Willebrand factor. *Blood* 86:1035, 1995
- Born GVR: Aggregation of blood platelets by adenosine diphosphate and its reversal. *Nature* 194:927, 1962

36. Poole AW, Watson SP: Regulation of cytosolic calcium by collagen in single human platelets. *Br J Pharmacol* 115:101, 1995
37. Wang X, Yanagi S, Yang C, Inatome R, Yamamura H: Tyrosine phosphorylation and Syk activation are involved in thrombin-induced aggregation of epinephrine-potentiated platelets. *J Biochem* 121:325, 1997
38. Blake RA, Schieven GL, Watson SP: Collagen stimulates tyrosine phosphorylation of phospholipase C γ 2 but not phospholipase C γ 1 in human platelets. *FEBS Lett* 353:212, 1994
39. Yanaga F, Asselin J, Schieven GL, Watson SP: Phenylarsine oxide inhibits tyrosine phosphorylation of phospholipase C γ 2 in human platelets and phospholipase C γ 1 in NIH-3T3 fibroblasts. *FEBS Lett* 368:377, 1995
40. Briddon SJ, Watson SP: Evidence for the involvement of a Src-like kinase in signalling by collagen-related peptide in washed human platelets. *Br J Pharmacol* 122:102P, 1997
41. Ikeda Y, Handa M, Kamata T, Kawano K, Kawai Y, Watanabe K, Kawakami K, Sakai K, Fukuyama M, Itagaki I, Yoshioka A, Ruggeri ZM: Transmembrane calcium influx associated with von Willebrand factor binding to GP Ib in the initiation of shear-induced platelet aggregation. *Thromb Haemost* 69:496, 1993
42. Ruggeri ZM: von Willebrand factor. *J Clin Invest* 99:559, 1997
43. Oda A, Yokoyama K, Murata M, Tokuhira M, Nakamura K, Handa M, Watanabe K, Ikeda Y: Protein tyrosine phosphorylation in human platelets during shear stress-induced platelet aggregation (SIPA) is regulated by glycoprotein (GP) Ib/IX as well as GP IIb/IIIa and requires intact cytoskeleton and endogenous ADP. *Thromb Haemost* 74:736, 1995
44. Ozaki Y, Satoh K, Yatomi Y, Miura S, Fujimura Y, Kume S: Protein tyrosine phosphorylation in human platelets induced by interaction between glycoprotein Ib and von Willebrand factor. *Biochim Biophys Acta* 1243:482, 1995
45. Razdan K, Hellums JD, Kroll MH: Shear-stress-induced von Willebrand factor binding to platelets causes the activation of tyrosine kinase(s). *Biochem J* 302:681, 1994
46. Gao J, Zoller KE, Ginsberg MH, Brugge JS, Shattil SJ: Regulation of the pp72syk protein tyrosine kinase by platelet integrin $\alpha_{IIb}\beta_3$. *EMBO J* 16:6414, 1997
47. Taniguchi T, Kitagawa H, Yasue S, Yanagi S, Sakai K, Asahi M, Ohta S, Takeuchi F, Nakamura S, Yamamura H: Protein-tyrosine kinase p72syk is activated by thrombin and is negatively regulated through Ca $^{2+}$ mobilization in platelets. *J Biol Chem* 268:2277, 1993
48. Reth M: Antigen receptor tail clue. *Nature* 338:383, 1989
49. Watson SP, Gibbins J: Collagen receptor signalling in platelets: Extending the role of the ITAM. *Immunol Today* 19:260, 1998
50. Ichinohe T, Takayama H, Ezumi Y, Yanagi S, Yamamura H, Okuma M: Cyclic AMP-insensitive activation of c-Src and Syk protein-tyrosine kinases through platelet membrane glycoprotein VI. *J Biol Chem* 270:28029, 1995
51. Sada K, Yanagi S, Yamamura H: Activation of p72syk by thrombin in a cell-free system. *Biochem Biophys Res Commun* 200:1, 1994
52. Cheng AM, Rowley B, Pao W, Hayday A, Bolen JB, Pawson T: Syk tyrosine kinase required for mouse viability and B-cell development. *Nature* 378:303, 1995
53. Turner M, Mee PJ, Costello PS, Williams O, Price AA, Duddy LP, Furlong MT, Geahlen RL, Tybulewicz VL: Perinatal lethality and blocked B-cell development in mice lacking the tyrosine kinase Syk. *Nature* 378:298, 1995
54. Ravetch JV, Kinetic JP: Fc receptors. *Annu Rev Immunol* 9:457, 1991
55. Amoui M, Draber P, Draberova L: Src-family selective tyrosine kinase inhibitor, PPI1, inhibits both Fc ϵ RI- and Thy-1-mediated activation of rat basophilic leukaemia cells. *Eur J Immunol* 27:1881, 1997
56. Daeron M: Fc receptor biology. *Annu Rev Immunol* 15:203, 1997
57. Stenberg PE, Pestina TI, Barrie RJ, Jackson CW: The Src family kinases, Fgr, Fyn, Lck and Lyn, colocalise with coated membranes in platelets. *Blood* 89:2384, 1997
58. Zhao Y-H, Krueger JG, Sudol M: Expression of cellular-yes protein in mammalian tissues. *Oncogene* 5:1629, 1990
59. Horak ID, Corcoran ML, Thompson PA, Wahl LM, Bolen JB: Expression of pp60fyn in human platelets. *Oncogene* 5:597, 1990
60. Golden A, Nemeth SP, Brugge JS: Blood platelets express high levels of the pp60c-src-specific tyrosine kinase activity. *Proc Natl Acad Sci USA* 83:852, 1986
61. Jackson SP, Schoenwaelder SM, Yuan Y, Rabinowitz I, Salem HH, Mitchell CA: Adhesion receptor activation of phosphatidylinositol 3-kinase. von Willebrand factor stimulates the cytoskeletal association and activation of phosphatidylinositol 3-kinase and pp60c-src in human platelets. *J Biol Chem* 269:27093, 1994
62. Duchemin AM, Anderson CL: Association of non-receptor protein tyrosine kinases with the Fc gamma RI/gamma-chain complex in monocytic cells. *J Immunol* 158:865, 1997
63. Keely PJ, Parise LV: The $\alpha_2\beta_1$ integrin is a necessary co-receptor for collagen-induced activation of Syk and the subsequent phosphorylation of phospholipase C γ 2 in platelets. *J Biol Chem* 271:26668, 1996
64. Dibirdik I, Kristupaitis D, Kurosaki T, Tuel-Ahlgren L, Chu A, Pond D, Tuong D, Luben R, Uckun FM: Stimulation of Src family protein-tyrosine kinases as a proximal and mandatory step for Syk kinase-dependent phospholipase C γ 2 activation in lymphoma B cells exposed to low energy electromagnetic fields. *J Biol Chem* 273:4035, 1998
65. Law CL, Chandran KA, Sidorenko SP, Clark EA: Phospholipase C γ 1 interacts with conserved phosphotyrosyl residues in the linker region of Syk and is a substrate for Syk. *Mol Cell Biol* 16:1305, 1996
66. Hanke JH, Gardner JP, Dow RL, Changelian PS, Brissette WH, Weringer EJ, Pollok BA, Connelly PA: Discovery of a novel, potent, and Src-family-selective tyrosine kinase inhibitor. Study of Lck- and Fyn-dependent T cell activation. *J Biol Chem* 271:695, 1996
67. Falasca M, Logan SK, Lehto VP, Baccante G, Lammon MA, Schlessinger J: Activation of phospholipase C γ by PI 3-kinase-induced PH domain-mediated membrane targeting. *EMBO J* 17:414, 1998
68. Rhee SG, Bae YS: Regulation of phosphoinositide-specific phospholipase C isozymes. *J Biol Chem* 272:15045, 1997
69. Gratacap M, Payrastra B, Viala C, Mauco G, Plantavid M, Chap H: Phosphatidylinositol 3,4,5-triphosphate-dependent stimulation of phospholipase C γ 2 is an early key event in Fc γ RIIA-mediated activation of human platelets. *J Biol Chem* 273:24314, 1998
70. Rameh LE, Rhee SG, Spokes K, Kazlauskas A, Cantley LC, Cantley LG: Phosphoinositide 3-kinase regulates phospholipase C γ -mediated calcium signaling. *J Biol Chem* 273:23750, 1998
71. Bae YS, Cantley LG, Chen CS, Kim SR, Kwon KS, Rhee SG: Activation of phospholipase C γ by phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem* 273:4465, 1998
72. Andrews RK, Harris SJ, McNally T, Berndt MC: Binding of purified 14-3-3 zeta signaling protein to discrete amino acid sequences within the cytoplasmic domain of the platelet membrane glycoprotein Ib-IX-V complex. *Biochemistry* 37:638, 1998
73. Calverley DC, Kavanagh TJ, Roth GJ: Human signaling protein 14-3-3zeta interacts with platelet glycoprotein Ib subunits Ibalpha and Ibbeta. *Blood* 91:1295, 1998
74. Du X, Harris SJ, Tetaz TJ, Ginsberg MH, Berndt MC: Association of a phospholipase A2 (14-3-3 protein) with the platelet glycoprotein Ib-IX complex. *J Biol Chem* 269:18287, 1994
75. Du X, Fox JE, Pei S: Identification of a binding sequence for the 14-3-3 protein within the cytoplasmic domain of the adhesion receptor, platelet glycoprotein Ib alpha. *J Biol Chem* 271:7362, 1996



blood[®]

1999 94: 1648-1656

Glycoprotein Ib-V-IX, a Receptor for von Willebrand Factor, Couples Physically and Functionally to the Fc Receptor γ -Chain, Fyn, and Lyn to Activate Human Platelets

Shahrokh Falati, Christine E. Edmead and Alastair W. Poole

Updated information and services can be found at:
<http://www.bloodjournal.org/content/94/5/1648.full.html>

Articles on similar topics can be found in the following Blood collections
[Hemostasis, Thrombosis, and Vascular Biology](#) (2485 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>